

# PROJECTIONS OF PATTERNS AND MODE INTERACTIONS

November 6, 2017

SOFIA B.S.D. CASTRO, ISABEL S. LABOURIAU, JULIANE F. OLIVEIRA

ABSTRACT. We study solutions of bifurcation problems with periodic boundary conditions, with periods in an  $n + 1$ -dimensional lattice and their projection into  $n$ -dimensional space through integration of the last variable. We show that generically the projection of a single mode solution is a mode interaction. This can be applied to the study of black-eye patterns.

## 1. INTRODUCTION

The aim of this article is to contribute to the understanding of models for the formation of periodic patterns in a thin layer. Experiments in thin layers are often modelled as 2-dimensional rather than 3-dimensional problems. The observation of results in such experiments is achieved through a 2-dimensional medium which is the top layer. In such instances, the observed patterns are 2-dimensional projections of the 3-dimensional objects in the thin layer.

An early example consists of the results found in reaction-diffusion experiments in the Turing instability regime, [24]. Typically, a reaction occurs in a thin layer of gel, fed by diffusion from one or two faces with chemicals contained in stirred tanks. The first reaction that provided Turing instabilities was in experiments on the chlorite-iodide-malonic-acid (CIMA) reaction, [3]. The pattern itself and its observed state can occur in different dimensions, see [15, 20, 12]. This happens for instance when an experiment is done in a 3-dimensional medium but the patterns are only observed on its surface, a 2-dimensional object, see [15, Section 4]. The interpretation of this 2-dimensional outcome is subject to discussion: the black-eye pattern observed by [20] has been explained both as a mode interaction in 2 dimensions in [13, 27], and described as a projection of a fully 3-dimensional pattern in [12]. The main challenge is to choose one of these descriptions. Our results in Section 5 indicate that these descriptions may coincide. We show that often the projection of a 3-dimensional single mode (the explanation provided in [12]) bifurcation is a 2-dimensional mode interaction (the explanation supported by [13, 27]). The comments of [27] concerning the monolayers that do or do not support the occurrence and observation of a black-eye pattern can be related to the width of the projection band (defined in Subsection 2.1).

Projection has proved to be a good way to model experiments. What is believed to be the first evidence of projection on the CIMA reaction can be found in [26, Chapter 13]. The author gives details about the geometry of the formation of wave patterns in malonic-acid reaction performed on sufficiently thin layers. In [15] the authors conducted experiments on the CIMA reaction and aimed at describing experimental observations of spontaneous symmetry breaking phenomena associated with steady-state instabilities. In [15, Section 4], the authors highlight the natural environment we must consider when we carry out CIMA reactions, in particular they state that all of their observations were based on projection of 3-dimensional structures. Moreover, the regions where Turing patterns are observed are associated by projection to a body-centred cubic lattice. More discussion on this can be found in [9, 16, 2, 23]. As reaction and diffusion progress bifurcations may occur. Bifurcating solutions are detected by observation of the top, 2-dimensional, layer. This poses the question of how solutions of a 3-dimensional object appear in their 2-dimensional observation.

Although we are motivated by the 2-dimensional projection of 3-dimensional objects, we study the relation between patterns in a  $(n + 1)$ -dimensional space and their projection in a  $n$ -dimensional space. Since a pattern consists of the level curves of periodic functions, we characterise the space of projected

periodic functions, describe the projection of orbits in the dual lattice and provide a decomposition of the span of projected irreducible subspaces. We achieve this by an algebraic approach to the relevant symmetry-related objects that contribute to the description of patterns. After some preliminary explanations, the main results are stated at the end of Section 2.

Information about which symmetries are preserved under projection has been obtained by Labouriau and Pinho in [17, 21]. In Oliveira *et al.* [19], we describe which symmetries can lead to a projected function with hexagonal symmetry. In studying the dynamics of  $(n+1)$ -dimensional patterns by observing their  $n$ -dimensional projection, it is desirable to extract information from the symmetries that can be seen in the  $n$ -dimensional space directly. The present article contributes to this point.

## 2. BIFURCATION PROBLEMS WITH EUCLIDEAN SYMMETRY

This section contains a rigorous formulation of the setting of the article and a statement of its main results. For this, definitions and basic results that are used in the article are stated here. More information on crystallographic groups may be found in Miller [18], Armstrong [1, chapters 24 to 26], Senechal [22], and the International Tables for Crystallography (ITC) volume A [14]. Results on equivariant bifurcation theory can be found in Golubitsky and Stewart's book [10]. For mode interactions we refer the reader to Castro [4, 5, 6] and [11, Ch XIX and XX].

Let  $E(n+1)$  be the *Euclidean group* of all isometries on  $\mathbb{R}^{n+1}$ , that may be described as the semi-direct sum  $E(n+1) \cong \mathbb{R}^{n+1} \dot{+} O(n+1)$ . Its elements are ordered pairs,  $(v, \delta)$ , where  $v \in \mathbb{R}^{n+1}$  is a translation and  $\delta$  is an element of the orthogonal group  $O(n+1)$ . The group operation is  $(v_1, \delta_1) \cdot (v_2, \delta_2) = (v_1 + \delta_1 v_2, \delta_1 \delta_2)$ . Elements  $(v, \delta) \in E(n+1)$  act on functions  $f: \mathbb{R}^{n+1} \rightarrow \mathbb{R}$  by  $((v, \delta) \cdot f)(x) = f(\delta^{-1}(x - v))$ .

Consider a one parameter family of partial differential equations

$$(1) \quad \frac{\partial u}{\partial t}(x, t) = \mathcal{F}(u(x, t), \lambda)$$

where  $\mathcal{F}: \mathcal{X} \times \mathbb{R} \rightarrow \mathcal{Y}$  is an operator between suitable function spaces  $\mathcal{X}$  and  $\mathcal{Y}$  and  $\lambda \in \mathbb{R}$  is a bifurcation parameter. The function  $u: \mathbb{R}^{n+1} \times \mathbb{R} \rightarrow \mathbb{R}$  in  $\mathcal{X}$  is a function of a spatial variable  $x \in \mathbb{R}^{n+1}$  and of time  $t$ .

Suppose that  $\mathcal{F}$  is *equivariant* under the Euclidean group  $E(n+1)$ , that is,

$$\gamma \cdot \mathcal{F}(u, \lambda) = \mathcal{F}(\gamma \cdot u, \lambda), \quad \text{for all } \gamma \in E(n+1).$$

Equilibria of (1) are time independent solutions  $u(x)$  that satisfy  $\mathcal{F}(u(x), \lambda) = 0$ . They are also called steady-states.

We give a brief description of a standard method to use symmetries to study the way steady-states in (1) bifurcate from the trivial solution  $\mathcal{F}(0, \lambda) \equiv 0$ . Details may be found in [8, 10]. The first step is to consider an  $(n+1)$ -dimensional lattice,  $\mathcal{L} \subset \mathbb{R}^{n+1}$ , a set generated over the integers by  $n+1$  linearly independent elements  $l_1, \dots, l_{n+1} \in \mathbb{R}^{n+1}$ , and to restrict the problem to a space  $\mathcal{X}_{\mathcal{L}}$  of functions periodic under  $\mathcal{L}$ .

There are natural symmetries in this space that we aim to explore to simplify the problem by applying the theory of bifurcation with symmetry. The restricted problem is equivariant under the action of the group  $\Gamma_{\mathcal{L}}$ , the largest group constructed from  $E(n+1)$  that leaves the space  $\mathcal{X}_{\mathcal{L}}$  invariant. Translations map  $\mathcal{X}_{\mathcal{L}}$  into itself, the action of translations on  $\mathcal{X}_{\mathcal{L}}$  is that of the torus  $\mathbb{T}^{n+1} = \mathbb{R}^{n+1}/\mathcal{L}$ . The subgroup of  $O(n+1)$  that maps  $\mathcal{L}$  to itself, and thus leaves  $\mathcal{X}_{\mathcal{L}}$  invariant, forms the *holohedry* of  $\mathcal{L}$  (called *point symmetry of the lattice* in [14]) and is denoted by  $H_{\mathcal{L}}$ . The holohedry is always finite, see [22, Sec. 2.4.2]. The group  $\Gamma_{\mathcal{L}}$  is thus compact and can be written as a semi-direct sum

$$\Gamma_{\mathcal{L}} = \mathbb{R}^{n+1}/\mathcal{L} \dot{+} H_{\mathcal{L}} = \mathbb{T}^{n+1} \dot{+} H_{\mathcal{L}}.$$

The next step is to analyse the bifurcation of solutions to the restricted problem. A steady-state bifurcation at  $\lambda = 0$  occurs when the linearisation of  $\mathcal{F}(0, 0)$  has a non-trivial kernel  $V \subset \mathcal{X}_{\mathcal{L}}$ . The kernel is always  $\Gamma_{\mathcal{L}}$ -invariant in the sense that if  $f \in V$  then  $\gamma \cdot f \in V$  for all  $\gamma \in \Gamma_{\mathcal{L}}$ . Since  $\Gamma_{\mathcal{L}}$  is compact, we expect  $V$  to be finite dimensional. This simplifies the problem considerably, as the study of bifurcating solutions is reduced to a  $\Gamma_{\mathcal{L}}$ -equivariant bifurcation problem defined in a finite dimensional space  $V$ . It also allows the use of equivariant bifurcation theory, where solution branches can then be obtained

using the Equivariant Branching Lemma [10, Lemma 1.31], independently proved by Cicogna [7] and Vanderbauwhede [25].

Next we provide some elementary definitions. Let  $\Gamma$  be a symmetry group of isometries acting on a vector space  $V$ . The *orbit* of  $v \in V$  under this action is the set  $\Gamma \cdot v = \{\gamma v : \gamma \in \Gamma\}$ . The *isotropy subgroup*  $\Sigma_v$  of  $v \in V$  is  $\Sigma_v = \{\gamma \in \Gamma : \gamma v = v\}$ . We say that a vector subspace  $W \subset V$  is  $\Gamma$ -*invariant* if  $\gamma \in \Gamma$  and  $v \in W$  implies  $\gamma v \in W$ . A  $\Gamma$ -invariant subspace  $W \subset V$  is  $\Gamma$ -*irreducible* if  $W$  contains no proper non-trivial  $\Gamma$ -invariant subspace. In this case, a problem defined in  $W$  cannot be reduced further.

Consider a  $\Gamma$ -equivariant bifurcation problem :

$$(2) \quad \frac{dz}{dt} = g(z, \lambda) \quad \text{with} \quad g(0, \lambda) = 0$$

with  $g : V \times \mathbb{R} \rightarrow V$  satisfying  $g(\gamma z, \lambda) = \gamma g(z, \lambda)$  for all  $\gamma \in \Gamma$  and all  $(z, \lambda) \in V \times \mathbb{R}$ .

**Definition 1.** The bifurcation problem (2) has:

- a *single mode bifurcation* at  $\lambda = 0$  if the kernel of  $(dg)_{0,0}$  is non-trivial and  $\Gamma$ -irreducible;
- an *r-mode interaction* at  $\lambda = 0$  if the kernel of  $(dg)_{0,0}$  can be decomposed as the direct sum of  $r > 1$  non-trivial components that are  $\Gamma$ -irreducible.

A characterisation of  $\Gamma_{\mathcal{L}}$ -irreducible subspaces of  $\mathcal{X}_{\mathcal{L}}$  is given by Dionne and Golubitsky [8], as follows.

Let  $\langle \cdot, \cdot \rangle$  be the usual inner product in  $\mathbb{R}^{n+1}$ . We assume that all the functions in  $\mathcal{X}_{\mathcal{L}}$  admit a Fourier expansion in terms of the *waves*

$$\omega_k(x, y) = \exp(2\pi i \langle k, (x, y) \rangle),$$

where  $k$  is a *wave vector* in the dual lattice,  $\mathcal{L}^* = \{k \in \mathbb{R}^{n+1}; \langle k, l_i \rangle \in \mathbb{Z}, i = 1, \dots, n+1\}$ , of  $\mathcal{L}$  and with the notation  $(x, y) \in \mathbb{R}^{n+1}$ ,  $x \in \mathbb{R}^n$ ,  $y \in \mathbb{R}$ .

Subspaces of  $\mathcal{X}_{\mathcal{L}}$  that are  $\Gamma_{\mathcal{L}}$ -irreducible must be, in particular,  $\mathbb{T}^{n+1}$ -invariant. Given  $\ell \in \mathbb{T}^{n+1}$  and  $k \in \mathcal{L}^*$  we have  $\ell \cdot \omega_k(x, y) = \omega_k(-\ell)\omega_k(x, y)$ . Hence, the two-dimensional subspaces

$$(3) \quad V_k = \{Re(z\omega_k(x, y)); z \in \mathbb{C}\}$$

are  $\mathbb{T}^{n+1}$ -irreducible. Moreover,  $V_k$  and  $V_{k'}$  are distinct  $\mathbb{T}^{n+1}$  representations if  $k \neq \pm k'$ .

The action of  $\gamma \in H_{\mathcal{L}}$  on  $\omega_k(x, y)$  satisfies

$$(4) \quad \gamma \omega_k(x, y) = \exp(2\pi i \langle k, \gamma^{-1}(x, y) \rangle) = \exp(2\pi i \langle \gamma k, (x, y) \rangle) = \omega_{\gamma k}(x, y)$$

by the orthogonality of  $\gamma$ . Then, we have:

**Proposition 1** (Dionne and Golubitsky [8]). *The space  $V = V_{k_1} \oplus \dots \oplus V_{k_s} \subset \mathcal{L}$  is  $\Gamma_{\mathcal{L}}$ -irreducible if and only if  $\{\pm k_1, \dots, \pm k_s\}$  is an  $H_{\mathcal{L}}$ -orbit in  $\mathcal{L}^*$ . In particular,  $2s$  divides the order of  $H_{\mathcal{L}}$ .*

We say that  $V$  is *generated by the orbit* of  $k$ . Since  $H_{\mathcal{L}} \subset O(n+1)$ , it follows that all the  $k_j$  ( $j = 1, \dots, s$ ) have the same norm.

## 2.1. Projections of $\mathcal{L}$ -periodic Functions.

**Definition 2.** For  $f \in \mathcal{X}_{\mathcal{L}}$  and  $y_0 > 0$ , the *projection operator*  $\Pi_{y_0}$  is given by:

$$\Pi_{y_0}(f)(x) = \int_0^{y_0} f(x, y) dy.$$

The region  $\{(x, y) \in \mathbb{R}^{n+1} : 0 \leq y \leq y_0\}$  is called the *band of projection*.

Hence, we are interested in describing the effect of the projection  $\Pi_{y_0}$  on  $\Gamma_{\mathcal{L}}$ -irreducible subspaces of  $\mathcal{X}_{\mathcal{L}}$ .

Pinho and Labouriau [17, 21] have characterised the group of symmetries of projected functions in the general case of functions invariant under a crystallographic group. Applying [17, Proposition 3.1], for  $n = 1$ , or [21, Theorem 1.2], for general  $n$ , to the special case where the crystallographic group only contains translations, it follows:

**Proposition 2.** *All functions in  $\Pi_{y_0}(\mathcal{X}_{\mathcal{L}})$  are invariant under the action of the translation  $v \in \mathbb{R}^n$  if and only if one of the following conditions holds:*

- I)  $(v, 0) \in \mathcal{L}$ ;
- II)  $(0, y_0) \in \mathcal{L}$  and  $(v, y_1) \in \mathcal{L}$ , for some  $y_1 \in \mathbb{R}$ .

We denote by  $\Pi_{y_0}(\mathcal{L}) = \tilde{\mathcal{L}}$  the set of all translations under which the functions in  $\Pi_{y_0}(\mathcal{X}_{\mathcal{L}})$  are invariant.

The action of  $\Gamma_{\mathcal{L}}$  on  $\mathcal{X}_{\mathcal{L}}$  induces a similar action of a group on  $\Pi_{y_0}(\mathcal{X}_{\mathcal{L}})$ . Translations act as the torus  $\mathbb{R}^n/\tilde{\mathcal{L}}$ . To see this, let  $g = \Pi_{y_0}(f)$ . If  $(v, 0) \in \mathcal{L}$  then  $g(x - v) = \int_0^{y_0} f(x - v, y - 0) dy \in \Pi_{y_0}(\mathcal{X}_{\mathcal{L}})$ . If  $(0, y_0)$  and  $(v, y_1) \in \mathcal{L}$ , since  $f(x, y + y_0) = f(x, y)$  then,

$$\int_0^{y_0} f(x - v, y - y_1) dy = \int_{y_1}^{y_0+y_1} f(x - v, z) dz = g(x - v) \in \Pi_{y_0}(\mathcal{X}_{\mathcal{L}}).$$

Given  $\alpha \in O(n)$ , define  $\alpha_{\pm} \in O(n+1)$  by:

$$\alpha_+ := \begin{pmatrix} \alpha & 0 \\ 0 & 1 \end{pmatrix} \text{ and } \alpha_- := \begin{pmatrix} \alpha & 0 \\ 0 & -1 \end{pmatrix}.$$

If either  $\alpha_+$  or  $\alpha_- \in H_{\mathcal{L}}$ , then  $\alpha$  belongs to the holohedry of  $\tilde{\mathcal{L}}$ . For  $g = \Pi_{y_0}(f)$  we get  $\Pi_{y_0}(\alpha_+ \cdot f) = \alpha \cdot g$ . If, in addition  $(0, y_0) \in \mathcal{L}$ , then

$$\Pi_{y_0}(\alpha_- \cdot f)(x) = \int_0^{y_0} f(\alpha^{-1}x, -y) dy = \int_{-y_0}^0 f(\alpha^{-1}x, z) dz = \alpha \cdot g(x)$$

because we are integrating over a period. We will denote by  $\tilde{\mathcal{J}} = \Pi_{y_0}(H_{\mathcal{L}})$  the group of *induced orthogonal symmetries*. It follows that  $\alpha \in \tilde{\mathcal{J}}$  if and only if one of the conditions below holds:

- I)  $\alpha_+ \in H_{\mathcal{L}}$ ;
- II)  $(0, y_0) \in \mathcal{L}$  and  $\alpha_- \in H_{\mathcal{L}}$ .

**2.2. Outline of the article and informal statement of results.** Given a lattice  $\mathcal{L}$  and its projection  $\tilde{\mathcal{L}} = \Pi_{y_0}(\mathcal{L})$ , the aim of this article is to relate functions in the space  $\mathcal{X}_{\mathcal{L}}$  to their projections, that lie in the space  $\mathcal{X}_{\tilde{\mathcal{L}}}$ , in order to obtain information about projected patterns and possible mode interactions. The starting point is the following:

**1st result:** Given a lattice  $\mathcal{L}$  with  $\tilde{\mathcal{L}} = \Pi_{y_0}(\mathcal{L})$ , for almost all  $y_0 \in \mathbb{R}$  we have  $\Pi_{y_0}(\mathcal{X}_{\mathcal{L}}) = \mathcal{X}_{\tilde{\mathcal{L}}}$ .

This result is proved in Section 3, where we obtain explicitly necessary and sufficient conditions on  $y_0$  under which  $\Pi_{y_0}(\mathcal{X}_{\mathcal{L}}) = \mathcal{X}_{\tilde{\mathcal{L}}}$ . In such cases, all functions invariant under  $\tilde{\mathcal{L}}$  can be obtained by the projection of a function invariant under  $\mathcal{L}$ .

Since solutions whose existence is guaranteed by the Equivariant Branching Lemma have less symmetry than the whole group, the relation between the symmetry of a function and its projection helps clarify what possible symmetries appear in the projected observation. When the symmetry of a function has a correspondence in the projected lattice, we know which possible symmetries to expect in the observed projected solutions. The Equivariant Branching Lemma then states that generically the solutions with maximal projected symmetry are observed.

The main result concerns mode interactions:

**2nd result:** Given a lattice  $\mathcal{L}$  with  $\tilde{\mathcal{L}} = \Pi_{y_0}(\mathcal{L})$ , generically and for almost all  $y_0 \in \mathbb{R}$  the projection of a single mode is a mode interaction.

Explicit conditions for the genericity and restrictions on the values of  $y_0$  are given in Section 5, where this result is proved. This shows the compatibility of the explanations provided for the black-eye pattern by [13] and [12]: when the outcome of an experiment is observed through a projection, generically, what is originally a single mode solution is projected as an observed mode interaction. Hence a 3-dimensional single mode (as explained by [12]) may be perceived as a mode interaction (as interpreted by [13]).

Some algebraic results that are used in the proof are developed in Section 4. These results concern a description of the action of  $H_{\mathcal{L}}$  on the dual lattice  $\mathcal{L}^*$  and its behaviour under projection. The results of Sections 4 and 5 are illustrated by the projection from the simple cubic lattice in  $\mathbb{R}^3$ , in two different positions, as well as the projection of the body centred cubic lattice, in a position that creates black-eyes.

3. PROJECTION OF THE SPACE OF  $\mathcal{L}$ -PERIODIC FUNCTIONS

Given an  $(n+1)$ -dimensional lattice  $\mathcal{L}$  and its projection  $\tilde{\mathcal{L}} = \Pi_{y_0}(\mathcal{L})$ , the projection of the space  $\mathcal{X}_{\mathcal{L}}$  of all  $\mathcal{L}$ -periodic functions does not necessarily coincide with the space of all functions with period in  $\tilde{\mathcal{L}}$ . In this section we characterise the situation when this is true. We illustrate this with examples, the end of each one marked  $\bullet$ .

Denote by  $P: \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$  the projection  $P(x, y) = x$  and by  $\{y = 0\}$  the space  $\{(x, y) \in \mathbb{R}^{n+1}; y = 0\}$ .

**Lemma 3.** *Let  $\mathcal{L}$  be an  $(n+1)$ -dimensional lattice with  $\tilde{\mathcal{L}} = \Pi_{y_0}(\mathcal{L})$ . Then  $\tilde{\mathcal{L}}^* \subset P(\mathcal{L}^*)$ .*

*Proof.* We analyse the two cases of Proposition 2. If  $(0, y_0) \in \mathcal{L}$ , then  $\tilde{\mathcal{L}} = P(\mathcal{L})$ . If  $\tilde{k} \in \tilde{\mathcal{L}}^*$  and  $(v, z) \in \mathcal{L}$  then  $\langle \tilde{k}, v \rangle = \langle (\tilde{k}, z), (v, 0) \rangle \in \mathbb{Z}$  because  $v \in \tilde{\mathcal{L}}$  and hence  $(\tilde{k}, 0) \in \mathcal{L}^*$ , therefore  $\tilde{k} \in P(\mathcal{L}^*)$ , as we wanted.

If  $(0, y_0) \notin \mathcal{L}$  then  $\tilde{\mathcal{L}} = P(\mathcal{L} \cap \{y = 0\})$ . If  $(\tilde{k}, z) \in \mathcal{L}^*$  then for every  $v \in \tilde{\mathcal{L}}$  we have  $(v, 0) \in \mathcal{L}$  and thus  $\langle (\tilde{k}, z), (v, 0) \rangle = \langle \tilde{k}, v \rangle \in \mathbb{Z}$ , hence  $\tilde{k} \in \tilde{\mathcal{L}}^*$ , therefore  $\tilde{\mathcal{L}}^* = P(\mathcal{L}^*)$ .  $\square$

**Theorem 4.** *Let  $\mathcal{L} \subset \mathbb{R}^{n+1}$  be an  $(n+1)$ -dimensional lattice and  $\tilde{\mathcal{L}} = \Pi_{y_0}(\mathcal{L})$ . The space  $\Pi_{y_0}(\mathcal{X}_{\mathcal{L}})$  is equal to  $\mathcal{X}_{\tilde{\mathcal{L}}}$  if and only if for each  $\tilde{k} \in \tilde{\mathcal{L}}^*$  there exists  $z \in \mathbb{R}$  such that  $(\tilde{k}, z) \in \mathcal{L}^*$  and  $zy_0 \notin \mathbb{Z} \setminus \{0\}$ .*

*Proof.* Suppose that  $\Pi_{y_0}(\mathcal{X}_{\mathcal{L}}) = \mathcal{X}_{\tilde{\mathcal{L}}}$ . Then, for all  $\omega_{\tilde{k}} \in \mathcal{X}_{\tilde{\mathcal{L}}}$ , there exists a non-zero function  $f \in \mathcal{X}_{\mathcal{L}}$  such that

$$\omega_{\tilde{k}}(x) = \Pi_{y_0}(f)(x) \neq 0.$$

By Lemma 3 we know that  $\tilde{\mathcal{L}} \subset P(\mathcal{L})$ . Since  $f \in \mathcal{X}_{\mathcal{L}}$  admits a Fourier expansion in terms of the waves  $\omega_k$  then, without loss of generality we may take  $f = c \omega_{(\tilde{k}, z)}$ , for some  $(\tilde{k}, z) \in \mathcal{L}^*$ . Moreover since  $\Pi_{y_0}(f)(x) \neq 0$ , then  $\int_0^{y_0} w_z(y) dy \neq 0$ . Therefore,  $zy_0 \notin \mathbb{Z} \setminus \{0\}$ .

Conversely, it is clear from Proposition 2 that  $\Pi_{y_0}(\mathcal{X}_{\mathcal{L}}) \subseteq \mathcal{X}_{\tilde{\mathcal{L}}}$ . To get the other inclusion, we want to show that all the wave functions  $\omega_{\tilde{k}} \in \mathcal{X}_{\tilde{\mathcal{L}}}$  are in  $\Pi_{y_0}(\mathcal{X}_{\mathcal{L}})$ . Suppose that for each  $\tilde{k} \in \tilde{\mathcal{L}}^*$  there exists  $z \in \mathbb{R}$  such that  $(\tilde{k}, z) \in \mathcal{L}^*$  and  $zy_0 \notin \mathbb{Z} \setminus \{0\}$ . Then,

$$\Pi_{y_0}(\omega_{(\tilde{k}, z)}) = c_0 \omega_{\tilde{k}}, \quad \text{with } c_0 = \int_0^{y_0} w_z(y) dy.$$

Since  $zy_0 \notin \mathbb{Z} \setminus \{0\}$ , then  $c_0 \neq 0$ . Therefore  $\Pi_{y_0}(\mathcal{X}_{\mathcal{L}}) = \mathcal{X}_{\tilde{\mathcal{L}}}$ .  $\square$

From Theorem 4 it follows:

**Corollary 5.** *Let  $\mathcal{L} \subset \mathbb{R}^{n+1}$  be an  $(n+1)$ -dimensional lattice and  $\tilde{\mathcal{L}} = \Pi_{y_0}(\mathcal{L})$ . Then for almost all values of  $y_0$  we have  $\Pi_{y_0}(\mathcal{X}_{\mathcal{L}}) = \mathcal{X}_{\tilde{\mathcal{L}}}$ .*

*Proof.* By Lemma 3, for each  $\tilde{k} \in \tilde{\mathcal{L}}^*$  there exists  $z \in \mathbb{R}$  such that  $(\tilde{k}, z) \in \mathcal{L}^*$ . Let  $(k_1, k_2) \in \mathbb{R}^n \times \mathbb{R}$  and consider the discrete set

$$K = \bigcup_{\substack{k_2 \neq 0; \\ (k_1, k_2) \in \mathcal{L}^*}} \mathbb{Z} \cdot \frac{1}{k_2}$$

then the complement of  $K$  is a dense set in  $\mathbb{R}$ .

For  $y_0 \in \mathbb{R} \setminus K$ , we have  $\int_0^{y_0} w_{k_2}(y) dy \neq 0$ . This implies that, for all  $y_0 \notin K$ , we have  $\Pi_{y_0}(\mathcal{X}_{\mathcal{L}}) = \mathcal{X}_{\tilde{\mathcal{L}}}$ .  $\square$

 4. PROJECTION OF  $H_{\mathcal{L}}$ -ORBITS IN  $\mathcal{L}^*$ 

We want to describe the effect of  $\Pi_{y_0}$  on irreducible subspaces of  $\mathcal{X}_{\mathcal{L}}$ . Proposition 1 tells us that this is done by studying the orbit of elements  $k$  of  $\mathcal{L}^*$  under the holohedry  $H_{\mathcal{L}}$ . The projection of the irreducible subspace generated by this orbit can be decomposed into irreducible subspaces under the action of symmetries of  $\tilde{\mathcal{L}} = \Pi_{y_0}(\mathcal{L})$  that lie in the group  $\tilde{J} = \Pi_{y_0}(H_{\mathcal{L}})$  that was defined in Subsection 2.1 above. This is equivalent to decomposing  $P(H_{\mathcal{L}} \cdot k)$  into  $\tilde{J}$ -orbits in  $\tilde{\mathcal{L}}^*$ . To do this, we decompose  $H_{\mathcal{L}}$  into subsets such that the orbit of  $k$  under each of these subsets is projected into exactly one  $\tilde{J}$ -orbit.

Define the subset  $\tilde{J}^\uparrow$  of  $H_{\mathcal{L}}$  as

$$(5) \quad \tilde{J}^\uparrow = \{\alpha_+ \in H_{\mathcal{L}}; \alpha \in \tilde{J}\} \cup \{\alpha_- \in H_{\mathcal{L}}; \alpha \in \tilde{J}\}.$$




FIGURE 1. If  $P(\delta(k_1, k_2)) = k_1$ , then  $\delta(k_1, k_2)|_2 = \pm k_2$ . Conversely, if  $\delta(k_1, k_2)|_2 \neq \pm k_2$  then  $k_1$  and  $P(\delta(k_1, k_2))$  do not have the same norm and thus cannot be in the same  $\tilde{J}$ -orbit.

Then for every  $k = (\tilde{k}, z) \in \mathcal{L}^*$ , the projection  $P(\tilde{J}^\dagger \cdot k) \subset \tilde{J} \cdot \tilde{k}$ . However, the set of  $\delta \in H_{\mathcal{L}}$  such that  $P(\delta k) \in \tilde{J} \cdot \tilde{k}$  is in general larger than  $\tilde{J}^\dagger$ . Next, we describe the subset of  $\delta \in H_{\mathcal{L}}$  such that the projection by  $P$  of  $\delta k$  lies in  $\tilde{J} \cdot \tilde{k}$ .

Given  $k = (\tilde{k}, z) \in \mathcal{L}^*$  and  $\alpha \in \tilde{J}$ , it is convenient to define  $J_k^\alpha$ , the subset of  $H_{\mathcal{L}}$  given by:

$$J_k^\alpha = \{\delta \in H_{\mathcal{L}}; P(\delta(\tilde{k}, z)) = \alpha \tilde{k}\}.$$

Denote the union of the sets  $J_k^\alpha$  by

$$S_k = \bigcup_{\alpha \in \tilde{J}} J_k^\alpha.$$

Then the projection of  $S_k \cdot k$  satisfies  $P(S_k \cdot k) = \tilde{J} \cdot P(k)$ . We show in Theorem 7 that  $H_{\mathcal{L}} k$  is decomposed into orbits given by cosets  $\delta S_k$  and that  $P(\delta S_k \cdot k) = \tilde{J} \cdot P(k)$ . The main problem is that  $S_k$  is not necessarily a group. We start with some properties of the sets we have defined. From now on for  $v = (x, y) \in \mathbb{R}^{n+1}$ ,  $x \in \mathbb{R}^n$ ,  $y \in \mathbb{R}$  we use the notation  $v|_2 = y$ .

**Proposition 6.** *The following properties hold for  $k = (\tilde{k}, z) \in \mathcal{L}^*$ :*

- (i) *If  $\delta \in J_k^{I^{d_n}}$ , then the last coordinate,  $\delta k|_2$ , of  $\delta k$  is equal to  $\pm z$ .*
- (ii) *For  $\alpha \in \tilde{J}$  then  $J_k^\alpha = \gamma \cdot J_k^{I^{d_n}}$  where  $\gamma \in H_{\mathcal{L}}$  is either  $\gamma = \alpha_+$  or  $\gamma = \alpha_-$ .*
- (iii) *Let  $\Sigma_k$  be the isotropy subgroup of  $k$  in  $H_{\mathcal{L}}$ . Then either  $J_k^{I^{d_n}} = \Sigma_k$  or it is the disjoint union  $\Sigma_k \cup \beta_- \Sigma_k$ , for some  $\beta_- \in J_k^{I^{d_n}}$ .*
- (iv) *The set  $S_k$  satisfies  $S_k = \bigcup_{\alpha \in \tilde{J}} J_k^\alpha = \tilde{J}^\dagger J_k^{I^{d_n}}$ . Moreover,  $S_k \cdot k = \tilde{J}^\dagger \cdot k$ .*
- (v) *If  $J_k^{I^{d_n}}$  is a subgroup of  $H_{\mathcal{L}}$  and if  $\tilde{J}^\dagger J_k^{I^{d_n}} = J_k^{I^{d_n}} \tilde{J}^\dagger$ , then  $S_k$  is a subgroup of  $H_{\mathcal{L}}$ .*

*Proof.* For  $\delta \in J_k^{I^{d_n}}$ , we write  $\delta k = (\tilde{k}, (\delta k)|_2)$ . Then by orthogonality of  $H_{\mathcal{L}}$ :

$$\|(\tilde{k}, z)\| = \|\delta k\| = \|(\tilde{k}, \delta(\tilde{k}, z)|_2)\|$$

Therefore  $|(\delta k)|_2| = |z|$ . This proves item (i). See Figure 1 for an illustration of this property.

To prove item (ii), let  $\gamma = \alpha_+$  or  $\gamma = \alpha_-$ , depending on whether either  $\alpha_+$  or  $\alpha_-$  is in  $H_{\mathcal{L}}$ . Let  $\phi: J_k^{I^{d_n}} \rightarrow J_k^\alpha$  given by  $\delta \mapsto \phi(\delta) = \gamma \delta$ . We show that the map  $\phi$  is injective and onto.

In fact, if  $\phi(\delta_1) = \phi(\delta_2)$ , for some  $\delta_1, \delta_2 \in J_k^{I^{d_n}}$ , then  $\gamma \delta_1 = \gamma \delta_2$ , for  $\gamma = \alpha_+$  or  $\gamma = \alpha_-$ , implying that  $\delta_1 = \delta_2$ . Thus  $\phi$  is injective.

Now consider  $\rho \in J_k^\alpha$ , then there exist  $\alpha_\pm^{-1} \rho \in H_{\mathcal{L}}$  and  $\tilde{k}_2$ , such that

$$\alpha_\pm^{-1} \rho(\tilde{k}, z) = \alpha_\pm^{-1}(\alpha \tilde{k}, \tilde{k}_2) = (\tilde{k}, \pm \tilde{k}_2)$$

that is,  $\alpha_\pm^{-1} \rho \in J_{\tilde{k}}^{I^{d_n}}$  and  $\phi(\alpha_\pm^{-1} \rho) = \rho$ .

The items (iii) and (v) are now immediate.  $\square$

Note that item (iii) implies that, in general, if  $J_k^{I^{d_n}} \neq \Sigma_k$  then  $J_k^{I^{d_n}}$  is not a group, as the next example will show.

**Example 1.** Let  $\mathcal{L}_1$  be the simple 3-dimensional cubic lattice generated by  $(1, 0, 0)$ ,  $(0, 1, 0)$ ,  $(0, 0, 1)$ , with dual lattice  $\mathcal{L}_1^*$  generated by the same vectors.

The holohedry  $H_{\mathcal{L}_1}$  has 24 rotations, they are generated by: the identity  $Id_3$ ; three rotations of order 4:  $R_x$ : rotation about  $(1, 0, 0)$ ;  $R_y$ : rotation about  $(0, 1, 0)$  and  $R_z$ : rotation about  $(0, 0, 1)$ . The remaining rotations are denoted by  $R_v$  as a rotation about the axis  $v$ :  $R_{(1,1,1)}$ ,  $R_{(1,-1,-1)}$ ,  $R_{(1,-1,1)}$ ,  $R_{(1,1,-1)}$  of order 3, and  $R_{(1,0,1)}$ ,  $R_{(1,0,-1)}$ ,  $R_{(1,1,0)}$ ,  $R_{(1,-1,0)}$ ,  $R_{(0,1,1)}$  and  $R_{(0,1,-1)}$  of order 2.

The remaining elements of  $H_{\mathcal{L}_1}$  can be obtained by multiplying these matrices by  $-Id_3$ .

TABLE 1. Relation of the set  $S_k$  for all  $k \in \mathcal{L}_1^* \setminus \{(0, 0, 0)\}$  and the group  $H_{\mathcal{L}}$  for Example 1. Note that  $|S_{(a,b,b)}| = 32$ , hence  $S_{(a,b,b)}$  is not a subgroup of  $H_{\mathcal{L}}$ , since  $|H_{\mathcal{L}}| = 48$ .

$k$ with $abc \neq 0$ $a \neq b \neq c \neq a$	$\Sigma_k$	$J_k^{Id_2}$	$S_k$
$(a, 0, 0)$	$\Sigma_{(a,0,0)} = J_{(a,0,0)}^{Id_2}$	$J_{(a,0,0)}^{Id_2} = \{Id_3, R_x, R_x^2, R_x^3, -R_y^2, -R_z^2, -R_{(0,1,1)}, -R_{(0,1,-1)}\}$	$ S_{(a,0,0)}  = 32$
$(a, b, b)$	$\Sigma_{(a,b,b)} = \{Id_3, -R_{(0,1,-1)}\}$	$J_{(a,b,b)}^{Id_2} = \{Id_3, -R_z^2, R_x^3, -R_{(0,1,-1)}\}$	$ S_{(a,b,b)}  = 32$
$(a, a, b)$	$\Sigma_{(a,a,b)} = \{Id_3, -R_{(1,-1,0)}\}$	$J_{(a,a,b)}^{Id_2} = \{Id_3, -R_z^2, R_{(1,1,0)}, -R_{(1,-1,0)}\}$	$S_{(a,a,b)} = \tilde{\mathcal{J}}^\dagger$
$(a, a, 0)$	$\Sigma_{(a,a,0)} = J_{(a,a,0)}^{Id_2}$	$J_{(a,a,0)}^{Id_2} = \{Id_3, -R_z^2, R_{(1,1,0)}, -R_{(1,-1,0)}\}$	$S_{(a,a,0)} = \tilde{\mathcal{J}}^\dagger$
$(a, b, 0)$	$\Sigma_{(a,b,0)} = J_{(a,b,0)}^{Id_2}$	$J_{(a,b,0)}^{Id_2} = \{Id_3, -R_z^2\}$	$S_{(a,b,0)} = \tilde{\mathcal{J}}^\dagger$
$(a, b, c)$	$\Sigma_{(a,b,c)} = \{Id_3\}$	$J_{(a,b,c)}^{Id_2} = \{Id_3, -R_z^2\}$	$S_{(a,b,c)} = \tilde{\mathcal{J}}^\dagger$
$(a, a, a)$	$\Sigma_{(a,a,a)} = \{Id_3, R_{(1,1,1)}, R_{(1,1,1)}^2, -R_{(1,0,-1)}, -R_{(0,1,-1)}, -R_{(1,-1,0)}\}$	$J_{(a,a,a)}^{Id_2} = \Sigma_{(a,a,a)} \cup \{R_x^3, R_y, -R_z^2, -R_{(1,-1,-1)}, -R_{(1,-1,1)}, -R_{(1,1,0)}\}$	$S_{(a,a,a)} = H_{\mathcal{L}_1}$

For any  $y_0$ , the projected lattice  $\Pi_{y_0}(\mathcal{L}_1) = \tilde{\mathcal{L}}_1$  is generated by the vectors  $(1, 0)$ ,  $(0, 1)$ , its dual,  $\tilde{\mathcal{L}}_1^*$  is generated by the same vectors.

The subgroup  $\tilde{\mathcal{J}}^\dagger$  has order 16 and is given by  $\{\pm Id_3, \pm R_x^2, \pm R_y^2, \pm R_z, \pm R_z^2, \pm R_z^3, \pm R_{(1,1,0)}, \pm R_{(1,-1,0)}\}$ .

The orbit of an element  $(a, b, c) \in \mathcal{L}_1^*$  by  $H_{\mathcal{L}_1}$  is given by:

$$(6) \quad \{(\pm a, \pm b, \pm c), (\pm b, \pm a, \pm c)\} \cup \{(\pm a, \pm c, \pm b), (\pm c, \pm a, \pm b)\} \cup \{(\pm b, \pm c, \pm a), (\pm c, \pm b, \pm a)\}$$

and the projection  $P(H_{\mathcal{L}_1}(a, b, c))$  is given by:

$$\{(\pm a, \pm b), (\pm b, \pm a)\} \cup \{(\pm a, \pm c), (\pm c, \pm a)\} \cup \{(\pm c, \pm b), (\pm b, \pm c)\}$$

On the other hand, for any  $y_0 \in \mathbb{R}$ , the group  $\tilde{\mathcal{J}} = \Pi_{y_0}(H_{\mathcal{L}_1})$  is the dihedral group of symmetries of the square,  $D_4$ , generated by:

$$\gamma = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad \kappa = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

For every  $\alpha \in \tilde{\mathcal{J}}$  both  $\alpha_+$  and  $\alpha_- \in H_{\mathcal{L}}$ . The orbit  $\tilde{\mathcal{J}}(a, b)$  is  $\{(\pm a, \pm b), (\pm b, \pm a)\}$ .

In Table 1 we describe, for each  $k \in \mathcal{L}_1^* \setminus \{(0, 0, 0)\}$ , the subgroup  $\Sigma_k$  and the set  $J_k^{Id_2}$  as well as the way the set  $S_k$  and the group  $H_{\mathcal{L}}$  are related. Some special cases are worth mentioning. If  $b \neq 0 \neq a$  then  $J_{(a,b,b)}^{Id_2}$  is not a group, because it contains the order 4 element  $R_x^3$  but not its powers. In the other cases where  $J_k^{Id_2} \neq \Sigma_k$  we have that  $\beta_-$  has order 2 and  $J_k^{Id_2}$  is a group.

The only cases where  $S_k \neq \tilde{\mathcal{J}}$  are for  $k = (a, a, a)$ , where  $S_k$  is the whole group  $H_{\mathcal{L}_1}$  and for  $k = (a, b, b)$ ,  $0 \neq a \neq b$ . For the second case, note that  $\delta \in H_{\mathcal{L}}$  is not in  $S_{(a,b,b)}$  if  $\delta(a, b, b) = (\pm b, \pm b, \pm a)$ . There are 8 rotations with this property:  $R_y$  and  $R_y^3$ ; the order 2 rotations  $R_{(1,0,1)}$  and  $R_{(1,0,-1)}$ ; and one and only one rotation of order 3 around each of the four axes of order 3. Multiplying them by  $-Id_3$  yields 16 elements not in  $S_{(a,b,b)}$ , hence  $|S_{(a,b,b)}| = 32$  and thus  $S_{(a,b,b)}$  is not a subgroup of  $H_{\mathcal{L}_1}$ . •

**Example 2.** Consider now the lattice  $\mathcal{L}_2$  that has generators  $(1, 0, 0)$ ,  $(\frac{1}{2}, \frac{\sqrt{3}}{2}, 0)$ ,  $(\frac{1}{2}, \frac{\sqrt{3}}{6}, \frac{\sqrt{6}}{6})$  given by  $\mathcal{L}_2 = \frac{1}{\sqrt{2}}A\mathcal{L}_1$  where  $A$  is:

$$A = \begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} & 0 \\ \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{6}} & \frac{2}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{3}} \end{pmatrix}.$$

The lattice  $\mathcal{L}_2$  has holohedry  $H_{\mathcal{L}_2} = AH_{\mathcal{L}_1}A^{-1}$ .

If  $y_0 \neq n\sqrt{6}/2$  with  $n \in \mathbb{Z} \setminus \{0\}$ , then  $(0, 0, y_0) \notin \mathcal{L}_2$ . In this case  $\tilde{\mathcal{L}}_2$  is generated by  $(1, 0)$  and  $(\frac{1}{2}, \frac{\sqrt{3}}{2})$  and  $\tilde{\mathcal{J}}$  is isomorphic to the dihedral group  $D_3$ . However, the holohedry of  $\tilde{\mathcal{L}}_2$  is larger than  $\tilde{\mathcal{J}}$ . It contains rotations of order 6 and is isomorphic to  $D_6$ .

For  $y_0 = n\sqrt{6}/2$  with  $n \in \mathbb{Z} \setminus \{0\}$ , then  $\tilde{\mathcal{L}}_2$  is generated by  $(1, 0)$  and  $(\frac{1}{2}, \frac{\sqrt{3}}{6})$  and  $\tilde{\mathcal{J}}$  is isomorphic to the dihedral group  $D_6$ , coinciding with the holohedry of  $\tilde{\mathcal{L}}_2$ .

The dual lattice  $\mathcal{L}_2^* = \sqrt{2}A\mathcal{L}_1^*$  is generated by  $(0, 0, \sqrt{6})$ ,  $(1, -\frac{\sqrt{3}}{3}, -\frac{\sqrt{6}}{3})$ , and  $(0, \frac{2\sqrt{3}}{3}, -\frac{\sqrt{6}}{3})$ .

For  $k = (2, 0, 0) \in \mathcal{L}_2^*$ , the set  $J_{(2,0,0)}^{Id_2} = \Sigma_{(2,0,0)}$  is a subgroup of  $H_{\mathcal{L}_2}$ . The  $\tilde{\mathcal{J}}$ -orbit of  $P(2, 0, 0) = (2, 0)$  contains 6 elements, independently of  $y_0$ , and is

$$\tilde{\mathcal{J}} \cdot (2, 0) = \{(\pm 2, 0), (\pm 1, \pm\sqrt{3})\}.$$

We claim that  $S_{(2,0,0)}$  is not a group. To see this we compute

$$AR_yA^{-1} = \begin{pmatrix} \frac{1}{2} & \frac{\sqrt{3}}{6} & -\frac{2\sqrt{3}}{3} \\ -\frac{\sqrt{3}}{2} & \frac{1}{6} & -\frac{\sqrt{2}}{3} \\ 0 & \frac{2\sqrt{2}}{3} & \frac{1}{3} \end{pmatrix} \quad AR_y^2A^{-1} = \begin{pmatrix} 0 & -\frac{\sqrt{3}}{3} & -\frac{\sqrt{6}}{3} \\ -\frac{\sqrt{3}}{3} & -\frac{2}{3} & \frac{\sqrt{2}}{3} \\ -\frac{\sqrt{6}}{3} & \frac{\sqrt{2}}{3} & -\frac{1}{3} \end{pmatrix} \quad AR_y^3A^{-1} = \begin{pmatrix} \frac{1}{2} & -\frac{\sqrt{3}}{2} & 0 \\ \frac{\sqrt{3}}{6} & \frac{1}{6} & \frac{2\sqrt{2}}{3} \\ -\frac{\sqrt{6}}{3} & -\frac{\sqrt{2}}{3} & \frac{1}{3} \end{pmatrix}.$$

Then  $AR_yA^{-1} \in S_{(2,0,0)}$ , since  $P(AR_yA^{-1}(2, 0, 0)) = (1, -\sqrt{3})$  and yet the powers of  $AR_yA^{-1}$  are not in  $S_{(2,0,0)}$ , establishing the claim. •

We can now prove the main result of this section. See Figure 1.

**Theorem 7.** Consider an  $(n+1)$ -dimensional lattice  $\mathcal{L}$  with holohedry  $H_{\mathcal{L}}$ , and let  $\tilde{\mathcal{J}} = \Pi_{y_0}(H_{\mathcal{L}})$ . Let  $k \in \mathcal{L}^*$ . Then there are  $\delta_j \in H_{\mathcal{L}}$ ,  $j = 2, \dots, r$  yielding  $k_1 = k, k_2 = \delta_2 k, \dots, k_r = \delta_r k$  such that  $H_{\mathcal{L}} \cdot k$  is the disjoint union

$$H_{\mathcal{L}} \cdot k = (S_{k_1} \cdot k_1) \cup (S_{k_2} \cdot k_2) \cup \dots \cup (S_{k_r} \cdot k_r)$$

and therefore the projection  $P(H_{\mathcal{L}}k)$  is a disjoint union of  $\tilde{\mathcal{J}}$ -orbits.

*Proof.* Given  $\alpha \in \tilde{\mathcal{J}}$  let  $\alpha_* \in \tilde{\mathcal{J}}^\dagger \subset H_{\mathcal{L}}$  stand for either  $\alpha_+$  or  $\alpha_-$ , as appropriate. We claim that for any  $\delta \in H_{\mathcal{L}}$ , any  $\alpha \in \tilde{\mathcal{J}}$  and  $k \in \mathcal{L}^*$ , if  $\alpha_* \delta \in S_k$  then  $\delta \in S_k$ .

If  $\alpha_* \delta \in S_k$  then there exists  $\beta \in \tilde{\mathcal{J}}$  such that  $\alpha_* \delta k = \beta_* k$ , by items (i) and (ii) of Proposition 6. Hence,  $\delta k = \alpha_*^{-1} \beta_* k$  and since  $\alpha_*^{-1} \beta_* \in \tilde{\mathcal{J}}^\dagger$  this implies that  $\delta \in S_k$ , as claimed.

Now consider  $H_{\mathcal{L}} = S_k \cup S_k^c$  and  $\delta_2 \in S_k^c$ , where  $S_k^c$  is the complement of  $S_k$ . Then for any  $\alpha_* \in \tilde{\mathcal{J}}^\dagger$  we have  $\alpha_* \delta_2 \in S_k^c$ . In particular,  $\tilde{\mathcal{J}}^\dagger \cdot (\delta_2 k) \cap S_k \cdot k = \emptyset$ . On the other hand, by Proposition 6 item (iv), the set  $\tilde{\mathcal{J}}^\dagger \cdot (\delta_2 k) = S_{\delta_2 k} \cdot (\delta_2 k) \subset H_{\mathcal{L}} k$ . Thus, we can write

$$H_{\mathcal{L}} \cdot k = S_k \cdot k \cup S_{\delta_2 k} \cdot (\delta_2 k) \cup (S_k \cup S_{\delta_2 k} \delta_1)^c k.$$

Since  $H_{\mathcal{L}}$  is a finite group, we can repeat the process to obtain:

$$(7) \quad H_{\mathcal{L}} \cdot k = S_k \cdot k \cup S_{\delta_2 k} \cdot (\delta_2 k) \cup \dots \cup S_{\delta_r k} \cdot (\delta_r k).$$

Then, for  $u_1 = P(k)$  and  $u_i = P(\delta_i k)$ , we have the disjoint union

$$(8) \quad P(H_{\mathcal{L}} \cdot k) = \bigcup_{i=1}^r \tilde{\mathcal{J}} \cdot u_i$$

and the result follows.  $\square$

We are interested in the case when there are several components in (8). The next corollary provides a condition for this, in terms of the size of the different sets that are used in this section.

**Corollary 8.** *For  $k \in \mathcal{L}^*$ , the projection  $P(H_{\mathcal{L}} \cdot k)$  in (8) contains more than one  $\tilde{\mathcal{J}}$ -orbit if and only if*

$$(9) \quad \frac{|H_{\mathcal{L}}|}{|\tilde{\mathcal{J}}^\dagger|} > \frac{|\Sigma_k|}{|\tilde{\mathcal{J}}^\dagger \cap \Sigma_k|}.$$

*Proof.* We know that the cardinal number of the orbit of  $k$  by  $H_{\mathcal{L}}$  is the index

$$(10) \quad |H_{\mathcal{L}} k| = \frac{|H_{\mathcal{L}}|}{|\Sigma_k|}.$$

The elements of  $H_{\mathcal{L}} \cdot k$  that are projected into  $\tilde{\mathcal{J}} \cdot P(k)$  are those in  $S_k \cdot k$ , and this set coincides with  $\tilde{\mathcal{J}}^\dagger \cdot k$  by item (iv) of Proposition 6. Since the isotropy subgroup of  $k$  in  $\tilde{\mathcal{J}}^\dagger$  is  $\tilde{\mathcal{J}}^\dagger \cap \Sigma_k$ , then the number of different elements in  $H_{\mathcal{L}} \cdot k$  whose projection lies in  $\tilde{\mathcal{J}} \cdot P(k)$  is  $|\tilde{\mathcal{J}}^\dagger|/|\tilde{\mathcal{J}}^\dagger \cap \Sigma_k|$ . Therefore, there are points in  $H_{\mathcal{L}} \cdot k$  whose projection does not lie in  $\tilde{\mathcal{J}} \cdot P(k)$  if and only if

$$\frac{|H_{\mathcal{L}}|}{|\Sigma_k|} > \frac{|\tilde{\mathcal{J}}^\dagger|}{|\tilde{\mathcal{J}}^\dagger \cap \Sigma_k|}$$

and this condition is equivalent to (9).  $\square$

## 5. A $\Gamma_{\mathcal{L}}$ -IRREDUCIBLE DECOMPOSITION

We are ready to give a decomposition of the projection of  $\Gamma_{\mathcal{L}}$ -irreducible subspaces of  $\mathcal{X}_{\mathcal{L}}$ . As before we use the notation  $\tilde{\mathcal{J}} = \Pi_{y_0}(H_{\mathcal{L}})$ . The group  $\tilde{\Gamma}_{\tilde{\mathcal{L}}} = \mathbb{R}^n / \tilde{\mathcal{L}} + \tilde{\mathcal{J}}$  is contained in  $\Gamma_{\tilde{\mathcal{L}}}$ , but does not necessarily coincide with it, since the inclusion  $\tilde{\mathcal{J}} \subset H_{\tilde{\mathcal{L}}}$  may be strict. This is the case in Example 2 above, other examples appear in [19].

**Theorem 9.** *Let  $V$  be the  $\Gamma_{\mathcal{L}}$ -irreducible subspace of  $\mathcal{X}_{\mathcal{L}}$  generated by the orbit of  $k \in \mathcal{L}^*$ . The projection  $\Pi_{y_0}(V)$  is a  $\tilde{\Gamma}_{\tilde{\mathcal{L}}}$ -invariant subspace of  $\mathcal{X}_{\tilde{\mathcal{L}}}$ . If, moreover, condition (9) holds then  $\Pi_{y_0}(V)$  is the sum of more than one  $\tilde{\Gamma}_{\tilde{\mathcal{L}}}$ -irreducible subspace, for almost all values of  $y_0 \in \mathbb{R}$ .*

*Proof.* First note that  $\Pi_{y_0}(\omega_{(k_1, k_2)}) = c_0 \omega_{k_1}$  with  $c_0 = \int_0^{y_0} e^{2\pi i k_2 y} dy$  and thus,  $c_0 = 0$  if and only if  $k_2 y_0 \in \mathbb{Z} \setminus \{0\}$ . Therefore,  $\omega_{k_1}(x) \in \Pi_{y_0}(V)$  if and only if  $k_1 \in P(H_{\mathcal{L}} \cdot k \setminus Z_{y_0})$  where

$$Z_{y_0} = \{(x, n y_0) \in \mathbb{R}^{n+1} : n \in \mathbb{Z} \setminus \{0\}\}.$$

If  $H_{\mathcal{L}} \cdot k \subset Z_{y_0}$  then it follows that  $\Pi_{y_0}(V) = \{0\}$ , a subspace that is trivially  $\tilde{\Gamma}_{\tilde{\mathcal{L}}}$ -invariant.

TABLE 2. For the simple cubic lattice  $\mathcal{L}_1$  in Example 1, we know that  $|H_{\mathcal{L}}|/|\tilde{\mathcal{J}}^\dagger| = 3$  and for each  $k \in \mathcal{L}_1^*$  this table gives the numbers  $|\Sigma_k|$  and  $|\tilde{\mathcal{J}}^\dagger \cap \Sigma_k|$  and the number  $N$  of non-trivial  $\tilde{\mathcal{J}}$ -irreducible components of the projection of the  $\Gamma_{\mathcal{L}_1}$ -irreducible subspace  $V \in \mathcal{X}_{\mathcal{L}_1}$  generated by the orbit of  $k$ . Here  $a \neq b \neq c \neq a$  with  $a, b, c \in \mathbb{Z} \setminus \{0\}$ .

$k$	$(a, 0, 0)$	$(a, b, b)$	$(a, a, b)$	$(a, a, 0)$	$(a, b, 0)$	$(a, b, c)$	$(a, a, a)$
$ \Sigma_k $	8	2	2	4	2	1	6
$ \tilde{\mathcal{J}}^\dagger \cap \Sigma_k $	4	1	2	4	2	1	2
$N$	2	2	2	2	3	3	1

Suppose then that  $\hat{k} \in H_{\mathcal{L}} \cdot k \setminus Z_{y_0} \neq \emptyset$  and hence  $\omega_{k_1}(x) \in \Pi_{y_0}(V)$ . If  $\alpha \in \tilde{\mathcal{J}}$  then either  $\alpha_+$  or  $\alpha_- \in H_{\mathcal{L}}$ , let  $\alpha_*$  be the appropriate one. Then  $\alpha_* \hat{k} \notin Z_{y_0}$ , implying that  $\omega_{\alpha k_1}(x) \in \Pi_{y_0}(V)$ . Since from (4) we know that  $\alpha \omega_{k_1}(x) = \omega_{\alpha k_1}(x)$ , this shows that  $\Pi_{y_0}(V)$  is  $\tilde{\Gamma}_{\mathcal{L}}$ -invariant.

When condition (9) holds, by Corollary 8 we know that  $P(H_{\mathcal{L}} \cdot k)$  contains at least two distinct  $\tilde{\mathcal{J}}$ -orbits, say  $\tilde{\mathcal{J}} \cdot P(u_1, z_1)$  and  $\tilde{\mathcal{J}} \cdot P(u_2, z_2)$ . The set  $K = \frac{1}{z_1} \mathbb{Z} \cup \frac{1}{z_2} \mathbb{Z}$  is a discrete subset of  $\mathbb{R}$  and for all  $y_0 \in \mathbb{R}$  such that  $y_0 \notin K$ , the projection  $\Pi_{y_0}(V)$  has at least two irreducible components.  $\square$

**Corollary 10.** *Suppose that  $V$  is a  $\Gamma_{\mathcal{L}}$ -irreducible subspace of  $\mathcal{X}_{\mathcal{L}}$  generated by the orbit of  $k \in \mathcal{L}^*$ . If there exist  $(u_1, z_1)$  and  $(u_2, z_2)$  in  $H_{\mathcal{L}} \cdot k$  with  $z_1 \neq \pm z_2$ , then for almost all  $y_0 \in \mathbb{R}$  the projection of a single mode is a mode interaction.*

*Proof.* If  $z_1 \neq \pm z_2$ , then by item (ii) of Proposition 6, the  $\tilde{\mathcal{J}}$ -orbits of  $u_1$  and  $u_2$  are distinct, as in Figure 1. Since  $K = \frac{1}{z_1} \mathbb{Z} \cup \frac{1}{z_2} \mathbb{Z}$  is a discrete subset of  $\mathbb{R}$ , for all  $y_0 \notin K$  the projection  $\Pi_{y_0}(V)$  has at least two  $\tilde{\Gamma}_{\mathcal{L}}$ -irreducible components. Hence, we have an interaction of at least two modes.  $\square$

We return to the examples of Section 4.

**Example 1.** For the simple cubic lattice  $\mathcal{L}_1$ , it is always true that  $\tilde{\mathcal{L}}_1 = P(\tilde{\mathcal{L}} \cap \{y = 0\})$ , for any  $y_0$ . Hence, the projection of a  $\Gamma_{\mathcal{L}_1}$ -irreducible subspace of  $\mathcal{X}_{\mathcal{L}_1}$  is never the trivial subspace. The number of  $\tilde{\mathcal{J}}$ -irreducible components of  $\Pi_{y_0}(V)$ , where  $V$  is generated by  $k \in \mathcal{L}_1^*$ , is shown in Table 2 for each  $k \in \mathcal{L}_1^*$ . For instance, when  $k = (a, b, 0)$ ,  $a \neq b \in \mathbb{Z} \setminus \{0\}$  the projection  $\tilde{V} = \Pi_{y_0}(V)$  is the sum of three  $\tilde{\Gamma}_{\mathcal{L}_1}$ -irreducible subspaces, generated by the orbits of  $(a, 0)$ , of  $(0, b)$  and of  $(a, b)$ . For  $k = (a, 0, 0)$ ,  $a \neq 0$ , the projection  $\Pi_{y_0}(V)$  has the two irreducible components generated by the orbits of  $(a, 0)$  and of  $(0, 0)$ , one of the components consists of constant functions. The only case when the projection is irreducible is when  $V$  is generated by the orbit of  $k = (a, a, a)$ ,  $a \neq 0$ , where the only irreducible component is generated by the orbit of  $(a, a)$ .  $\bullet$

Example 2 is more typical, because the projection depends on  $y_0$ .

**Example 2.** The holohedry  $H_{\mathcal{L}_2}$  of the rotated cubic lattice  $\mathcal{L}_2$  has order 48. The subgroup  $\tilde{\mathcal{J}}^\dagger$  has order 12 if  $y_0 = n\sqrt{6}/2$  with  $n \in \mathbb{Z} \setminus \{0\}$ , otherwise it has order 6. Tables 3 and 4 give all the information necessary to apply Theorem 9 to this lattice. For each  $k$  there is at most a discrete set of values of  $y_0$  for which a single mode is projected into a single mode. Generically the projection is a mode interaction.  $\bullet$

**Example 3.** The body centered cubic lattice  $\mathcal{L}_3$  was used in [12] to obtain black-eye patterns by projection, like those of Figure 2. It is generated by  $(1/2, \sqrt{3}/2, 0)$ ,  $(1, 0, 0)$  and  $(1/2, \sqrt{3}/6, -\sqrt{6}/12)$ . It contains the lattice  $\mathcal{L}_2$  of Example 2 and the two lattices share the same holohedry. Generators for  $\mathcal{L}_3^* \subset \mathcal{L}_2^*$  are  $(1, \sqrt{3}, 0)$ ,  $(2, 0, 0)$  and  $(1, -\sqrt{3}/3, 2\sqrt{6}/3)$ . Since  $\mathcal{L}_3^*$  does not contain any vector of the form  $(a, \sqrt{3}a/3, \sqrt{6}a/3)$ , then the projection of every  $H_{\mathcal{L}_3}$ -orbit in  $\mathcal{L}_3^*$  contains more than one  $\tilde{\mathcal{J}}$ -orbit. For instance (see Figure 3), the orbit  $H_{\mathcal{L}_3} \cdot (1, \sqrt{3}, 0)$  has 12 elements:

$$\left\{ \pm(1, \sqrt{3}, 0), \pm(2, 0, 0), \pm(-1, \sqrt{3}, 0), \pm\left(1, \frac{\sqrt{3}}{3}, \frac{-2\sqrt{6}}{3}\right), \pm\left(0, \frac{2\sqrt{3}}{3}, \frac{2\sqrt{6}}{3}\right), \pm\left(-1, \frac{\sqrt{3}}{3}, \frac{-2\sqrt{6}}{3}\right) \right\}$$

TABLE 3. For the rotated cubic lattice  $\mathcal{L}_2$  in Example 2, and for  $y_0 = m\sqrt{6}/2$ ,  $m \in \mathbb{Z} \setminus \{0\}$ , we have that  $|H_{\mathcal{L}}|/|\tilde{\mathcal{J}}^\uparrow| = 4$ . For each  $k \in \mathcal{L}_1^*$  this table gives the transformed vector in  $\mathcal{L}_2^*$ , the numbers  $|\Sigma_k|$ , the set  $\tilde{\mathcal{J}}^\uparrow \cap \Sigma_k$ , the number  $|\Sigma_k|/|\Sigma_k \cap \tilde{\mathcal{J}}^\uparrow|$  and the restriction on  $y_0$ . If  $V$  is the irreducible subspace generated by the orbit of  $k$ , then the only case where  $\Pi_{y_0}(V)$  has only one irreducible component is  $k = \left(a, \frac{\sqrt{3}a}{3}, \frac{\sqrt{6}a}{3}\right)$ . A small change in  $y_0$  destroys this situation, see Table 4. Here  $a \neq b \neq c \neq a$  with  $a, b, c$  and  $n \in \mathbb{Z} \setminus \{0\}$ .

$k \in \mathcal{L}_1^*$	$\sqrt{2}Ak \in \mathcal{L}_2^*$	$ \Sigma_k $	$\Sigma_k \cap \tilde{\mathcal{J}}^\uparrow$	$ \Sigma_k / \Sigma_k \cap \tilde{\mathcal{J}}^\uparrow $	restriction
$(a, 0, 0)$	$\left(a, \frac{\sqrt{3}a}{3}, \frac{\sqrt{6}a}{3}\right)$	8	$\{Id_3, -R_{(0,1,1)}\}$	4	-
$(a, b, b)$	$\left(a - b, \sqrt{3}\left(\frac{a}{3} + b\right), \frac{\sqrt{6}a}{3}\right)$	2	$\mathbb{1}$	2	$y_0 \neq \frac{\sqrt{6}}{2a}n$
$(0, a, a)$	$(-a, \sqrt{3}a, 0)$	4	$\{Id_3, R_{(0,1,1)}\}$	2	-
$(a, b, 0)$	$\left(a - b, \frac{\sqrt{3}}{3}(a + b), \frac{\sqrt{6}}{3}(a + b)\right)$	2	$\mathbb{1}$	2	$y_0 \neq \frac{\sqrt{6}}{2(a+b)}n$
$(a, b, c)$	$\left(a - b, \frac{\sqrt{3}}{3}(a + b + 2c), \frac{\sqrt{6}}{3}(a + b - c)\right)$	1	$\mathbb{1}$	1	$y_0 \neq \frac{\sqrt{6}}{2(a+b-c)}n$
$(a, a, a)$	$\left(0, \frac{4\sqrt{3}a}{3}, \frac{\sqrt{6}a}{3}\right)$	6	$\{Id_3, R_{(1,-1,0)}\}$	3	$y_0 \neq \frac{\sqrt{6}}{2a}n$
$(a, b, a + b)$	$\left(a - b, \frac{2\sqrt{3}}{3}(a + b), 0\right)$	1	$\mathbb{1}$	1	-
$(a, 0, a)$	$\left(a, \frac{2\sqrt{3}a}{3}, 0\right)$	4	$\{Id_3, R_{(1,0,1)}\}$	2	-
$(a, a, 2a)$	$\left(0, \frac{4\sqrt{3}a}{3}, 0\right)$	2	$\{Id_3, -R_{(1,-1,0)}\}$	1	-

TABLE 4. For the rotated cubic lattice  $\mathcal{L}_2$  in Example 2, and  $y_0 \neq m\sqrt{6}/2$ ,  $m \in \mathbb{Z} \setminus \{0\}$ , we have that  $|H_{\mathcal{L}}|/|\tilde{\mathcal{J}}^\uparrow| = 8$ . For each  $k \in \mathcal{L}_1^*$  this table gives the transformed vector in  $\mathcal{L}_2^*$ , the numbers  $|\Sigma_k|$ , the set  $\tilde{\mathcal{J}}^\uparrow \cap \Sigma_k$ , the number  $|\Sigma_k|/|\Sigma_k \cap \tilde{\mathcal{J}}^\uparrow|$  and the restriction on  $y_0$ . If  $V$  is the irreducible subspace generated by the orbit of  $k$ , then the projection  $\Pi_{y_0}(V)$  has always more than one irreducible component as long as the restriction on  $y_0$  holds. Here  $a \neq b \neq c \neq a$  with  $a, b, c$  and  $n \in \mathbb{Z} \setminus \{0\}$ .

$k \in \mathcal{L}_1^*$	$\sqrt{2}Ak \in \mathcal{L}_2^*$	$ \Sigma_k $	$\Sigma_k \cap \tilde{\mathcal{J}}^\uparrow$	$ \Sigma_k / \Sigma_k \cap \tilde{\mathcal{J}}^\uparrow $	restriction
$(a, 0, 0)$	$\left(a, \frac{\sqrt{3}a}{3}, \frac{\sqrt{6}a}{3}\right)$	8	$\{Id_3, -R_{(0,1,1)}\}$	4	$y_0 \neq \frac{\sqrt{6}}{2a}n$
$(a, b, b)$	$\left(a - b, \sqrt{3}\left(\frac{a}{3} + b\right), \frac{\sqrt{6}a}{3}\right)$	2	$\mathbb{1}$	2	$y_0 \neq \frac{\sqrt{6}}{2a}n$
$(0, a, a)$	$(-a, \sqrt{3}a, 0)$	4	$\mathbb{1}$	4	-
$(a, b, 0)$	$\left(a - b, \frac{\sqrt{3}}{3}(a + b), \frac{\sqrt{6}}{3}(a + b)\right)$	2	$\mathbb{1}$	2	$y_0 \neq \frac{\sqrt{6}}{2(a+b)}n$
$(a, b, c)$	$\left(a - b, \frac{\sqrt{3}}{3}(a + b + 2c), \frac{\sqrt{6}}{3}(a + b - c)\right)$	1	$\mathbb{1}$	1	$y_0 \neq \frac{\sqrt{6}}{2(a+b-c)}n$
$(a, a, a)$	$\left(0, \frac{4\sqrt{3}a}{3}, \frac{\sqrt{6}a}{3}\right)$	6	$\mathbb{1}$	6	$y_0 \neq \frac{\sqrt{6}}{2a}n$
$(a, b, a + b)$	$\left(a - b, \frac{2\sqrt{3}}{3}(a + b), 0\right)$	1	$\mathbb{1}$	1	-
$(a, 0, a)$	$\left(a, \frac{2\sqrt{3}a}{3}, 0\right)$	4	$\mathbb{1}$	4	-
$(a, a, 2a)$	$\left(0, \frac{4\sqrt{3}a}{3}, 0\right)$	2	$\{Id_3, -R_{(1,-1,0)}\}$	1	-

and for  $y_0 = \sqrt{6}n/4$ ,  $n \in \mathbb{Z} \setminus \{0\}$  its projection consists of two  $\tilde{\mathcal{J}}$ -orbits, whereas for all other values of  $y_0$  it is formed by three  $\tilde{\mathcal{J}}$ -orbits.

An irreducible subspace  $V \subset \mathcal{X}_{\mathcal{L}_3}$  generated by the  $H_{\mathcal{L}_3}$ -orbit of  $k \in \mathcal{L}_3$  contains a maximally symmetric function

$$I_k(x, y, z) = \sum_{\delta \in H_{\mathcal{L}_3}} \omega_{\delta k}(x, y, z).$$




FIGURE 2. Black eye pattern, obtained projecting the function  $I_{(1/2, \sqrt{3}/2, 0)} \in \mathcal{X}_{\mathcal{L}_3}$  with a band of projection of width  $y_0 = \sqrt{6}/4$ .




FIGURE 3. The  $\tilde{\mathcal{J}}$ -orbits in the projection  $P(H_{\mathcal{L}_3}(1, \sqrt{3}, 0))$  in Example 3. For  $y_0 = \sqrt{6}n/12$ ,  $n \in \mathbb{Z} \setminus \{0\}$  the projection decomposes into two  $\tilde{\mathcal{J}}$ -orbits, shown as marked points in the two dotted circles. For other values of  $y_0$  the points with small norm in the projection lie in two different  $\tilde{\mathcal{J}}$ -orbits, indicated here by black or white filled points. This gives rise to a 3-mode interaction.

The subspace of  $V$  generated by  $I_k$  is the fixed-point subspace of the maximal subgroup  $H_{\mathcal{L}_3} \subset \Gamma_{\mathcal{L}_3}$ . The Equivariant Branching Lemma [10, Lemma 1.31] may be applied to obtain patterns bifurcating into this subspace. The pattern shown in Figure 2 is the projection  $\Pi_{y_0}(I_{(1/2, \sqrt{3}/2, 0)})$  for  $y_0 = \sqrt{6}/4$ .

For  $\tilde{k} \in \tilde{\mathcal{L}}_3$ , let  $\tilde{I}_{\tilde{k}}$  be given by

$$\tilde{I}_{\tilde{k}} = \sum_{\beta \in \tilde{\mathcal{J}}} \omega_{\beta \tilde{k}}(x, y).$$

For most values of  $y_0$ , the projected group  $\tilde{\mathcal{J}}$  is isomorphic to the dihedral group  $D_3$  of order 6. The  $H_{\mathcal{L}_3}$ -orbit of  $(1/2, \sqrt{3}/2, 0)$  is projected into three  $\tilde{\mathcal{J}}$ -orbits, shown in Figure 3. This means that  $\Pi_{y_0}(I_{(1/2, \sqrt{3}/2, 0)})$  is a linear combination of the three  $\tilde{\mathcal{J}}$ -invariant functions  $\tilde{I}_{(2,0)}$ ,  $\tilde{I}_{(1, \sqrt{3}/3)}$  and  $\tilde{I}_{(-1, -\sqrt{3}/3)}$ , where the first function is real and the third function is the complex conjugate of the second. The patterns created by these functions are shown in the upper part of Figure 4.

For  $y_0 = \sqrt{6}n/12$ ,  $n \in \mathbb{Z} \setminus \{0\}$ , the projected group  $\tilde{\mathcal{J}}$  is isomorphic to the dihedral group  $D_6$  of order 12. The  $H_{\mathcal{L}_3}$ -orbit of  $(1/2, \sqrt{3}/2, 0)$  is projected into two  $\tilde{\mathcal{J}}$ -orbits, since the  $\tilde{\mathcal{J}}$ -orbits of  $\pm(1, \sqrt{3}/3)$  coincide, as in Figure 3. Therefore  $\tilde{I}_{(1, \sqrt{3}/3)} = \tilde{I}_{(-1, -\sqrt{3}/3)}$ . This means that  $\Pi_{\sqrt{6}/4}(I_{(1/2, \sqrt{3}/2, 0)})$  is a linear combination of the two  $\tilde{\mathcal{J}}$ -invariant functions  $\tilde{I}_{(2,0)}$  and  $\tilde{I}_{(1, \sqrt{3}/3)}$ . For the geometrical consequences, see Figure 4. •

#### ACKNOWLEDGMENTS

The authors would like to thank Professors Miriam Manoel and Gabriela Gomes for useful comments and suggestions.

CMUP is supported by the European Regional Development Fund through the programme COMPETE and by the Portuguese Government through the Fundação para a Ciência e a Tecnologia (FCT) under the project PEst-C/MAT/UI0144/2011. J.F. Oliveira was supported by a grant from the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) of Brazil.




FIGURE 4. Top, left to right: level sets for  $\tilde{I}_{(2,0)} \in \mathbb{R}$ ,  $\text{Re}(\tilde{I}_{(1,\sqrt{3}/3)}) = \text{Re}(\tilde{I}_{(1,-\sqrt{3}/3)})$  and  $\text{Im}(\tilde{I}_{(1,\sqrt{3}/3)}) = -\text{Im}(\tilde{I}_{(1,-\sqrt{3}/3)})$  where  $\tilde{I}$  corresponds to  $y_0 \neq \sqrt{6}n/3$ ,  $n \in \mathbb{Z} \setminus \{0\}$ . Bottom: projections  $\Pi_{y_0} \left( I_{(1/2,\sqrt{3}/2,0)} \right)$  for different  $y_0$ , from left to right:  $y_0 = 2\sqrt{6}/3$ ,  $y_0 = \sqrt{6}/3$  and  $y_0 = 8\sqrt{6}/15$ . The pattern in the centre is the same as in Figure 2, a perfect black eye. Those on the sides are similar, but with less symmetry. Note that the patterns at the bottom are linear combinations of those on top, with different coefficients.

## REFERENCES

- [1] M. A. Armstrong, *Groups and Symmetry*, Springer-Verlag, 1988
- [2] P. Borckmans, G. Dewel, A. De Wit, E. Dulos, J. Boissonade, F. Gauffre and P. De Kepper, *Diffusive instabilities and chemical reactions*, International journal of bifurcation and chaos, **12** (2002) 2307–2332
- [3] V. Castets, E. Dulos, J. Boissonade and P. De Kepper, *Experimental evidence of a sustained standing Turing-type nonequilibrium chemical pattern*, Physical Review Letters, **64-24**(1990) 2953
- [4] S.B.S.D. Castro, *Mode Interactions with Symmetry*, PhD thesis University of Warwick, 1993
- [5] S.B.S.D. Castro, *Mode interactions with spherical symmetry*, International Journal of Bifurcation and Chaos **4-04** (1994) 885–904
- [6] S.B.S.D. Castro, *Mode interactions with symmetry*, Dynamics and Stability of Systems **10-1** (1995) 13–31
- [7] G. Cicogna, *Symmetry breakdown from bifurcations*, Lettere al Nuovo Cimento 31 (1981) 600-602.
- [8] B. Dionne and M. Golubitsky, *Planforms in two and three dimensions*, Z. Angew. Math. Phys.**43** (1992) 36–62
- [9] E. Dulos, P. Davies, B. Rudovics and P. De Kepper, *From quasi-2D to 3D Turing patterns in ramped systems*, Physica D Nonlinear Phenomena, **98** (1996) 53-66,
- [10] M. Golubitsky and I. Stewart, *The Symmetry Perspective* Birkhauser Verlag, 2002
- [11] M. Golubitsky, I.N. Stewart and D. Schaeffer, *Singularities and groups in bifurcation theory, volume II*, Springer-Verlag, 2012
- [12] M.G.M. Gomes, *Black-eye patterns: A representation of three-dimensional symmetries in thin domains*, Phys. Rev. E, **60** (1999) 3741–3747
- [13] H.G. Gunaratne, Q. Ouyang and H.L. Swinney, *Pattern formation in the presence of symmetries*, Phys. Rev. E, **50** (1994) 2802–2806
- [14] T. Hahn (ed.), *International Tables for Crystallography, 5th edition*, International Union of Crystallography, 2006
- [15] P. De Kepper, V. Castets, E. Dulos and J. Boissonade, *Turing-type chemical patterns in the chlorite-iodide-malonic acid reaction*, Physica D: Nonlinear Phenomena, **49** (1991) = 161–169
- [16] P. De Kepper, E. Dulos, J. Boissonade, A. De Wit, G. Dewel and P. Borckmans, *Reaction–diffusion patterns in confined chemical systems*, Journal of statistical physics, **101** (2000) 495–508

- [17] I.S. Labouriau and E.M. Pinho, *Symmetries of projected wallpaper patterns*, Mathematical Proceedings of the Cambridge Philosophical Society **141** (2006) 421–441
- [18] W. Miller, *Symmetry Groups and their Applications*, Academic Press, 1972
- [19] J.F. Oliveira, S.B.S.D. Castro and I.S. Labouriau, *Hexagonal projected symmetries*, Acta Crystallographica Section A **71** (2015) 549–558
- [20] Q. Ouyang and H.L. Swinney, *Onset and beyond Turing pattern formation*, In R. Kapral and K. Showalter (eds), *Chemical waves and patterns*, Kluwer, 1995, 269–295
- [21] E.M. Pinho and I.S. Labouriau, *On the Projection of Functions Invariant under the Action of a Crystallographic Group*, Journal of Pure and Applied Algebra **218** (2014) 37–51
- [22] M. Senechal, *Quasicrystals and geometry*, Cambridge University Press, 1996
- [23] I. Szalai, J. Horváth and P. De Kepper, *Contribution to an effective design method for stationary reaction-diffusion patterns*, Chaos: An Interdisciplinary Journal of Nonlinear Science, **25** (2015) 064311
- [24] A.M. Turing, *The chemical basis of morphogenesis*, Phil. Trans. R. Soc. London. Ser. B, **237** (1952) 37–72
- [25] A. Vanderbauwhede, *Local Bifurcation and Symmetry* Habilitation Thesis, Rijksuniversiteit Gent (1980) (also Res. Notes Math. 75. Pitman, Boston (1982)).
- [26] A.T. Winfree, *The geometry of biological time*, Springer-Verlag, 2001
- [27] C. Zhou, H. Guo and Q. Ouyang, *Experimental study of the dimensionality of lack-eye patterns*, Phys. Rev. E, **65** (2002) 036118

CENTRO DE MATEMÁTICA DA UNIVERSIDADE DO PORTO, RUA DO CAMPO ALEGRE 687, 4169-007, PORTO, PORTUGAL,  
AND J.F. OLIVEIRA, I.S.LABOURIAU — DEPARTAMENTO DE MATEMÁTICA, FACULDADE DE CIÊNCIAS DA UNIVERSIDADE DO  
PORTO, PORTUGAL, S.B.S.D.CASTRO — FACULDADE DE ECONOMIA, UNIVERSIDADE DO PORTO, RUA DR. ROBERTO FRIAS,  
4200-464, PORTO, PORTUGAL