

Primitive ideals and irreducible representations of $U_q(\mathfrak{sl}_4^+)$

Samuel A. Lopes *
Centro de Matemática da Universidade do Porto,
R. Campo Alegre 687, 4169-007 Porto, Portugal
E-mail: slopes@fc.up.pt

April 3, 2005

Abstract

Let $U_q(\mathfrak{g}^+)$ be the quantized enveloping algebra of the nilpotent Lie algebra $\mathfrak{g}^+ = \mathfrak{sl}_{n+1}^+$ which occurs as the positive part in the triangular decomposition of the simple Lie algebra \mathfrak{sl}_{n+1} of type A_n . Assuming the base field \mathbb{K} is algebraically closed and of characteristic 0, and that the parameter $q \in \mathbb{K}^*$ is not a root of unity, we define and study certain quotients of $U_q(\mathfrak{g}^+)$ which coincide with the Hayashi algebra when $n = 2$ (see [14], [2] and [13]). We show that these are simple Noetherian domains, with a trivial center and even Gelfand-Kirillov dimension. Hence, they are adequate quantum analogues of the Weyl algebras. In the remainder of the paper we study the primitive spectrum of $U_q(\mathfrak{sl}_4^+)$ in detail, much in the spirit of [15]. We determine all primitive ideals of $U_q(\mathfrak{sl}_4^+)$, compute their heights and find a simple $U_q(\mathfrak{sl}_4^+)$ -module corresponding to each primitive ideal of $U_q(\mathfrak{sl}_4^+)$.

1 Introduction

This paper is concerned with the primitive ideals of the quantized enveloping algebra $U_q(\mathfrak{g}^+)$ of the nilpotent Lie algebra $\mathfrak{g}^+ = \mathfrak{sl}_{n+1}^+$ of strictly upper triangular matrices of size $n + 1$, with an emphasis on $U_q(\mathfrak{sl}_4^+)$. In the classical case, the primitive factors of the enveloping algebra $U(\mathfrak{g}^+)$ of \mathfrak{g}^+ are isomorphic to Weyl algebras, and consequently the primitive ideals of $U(\mathfrak{g}^+)$ are simply its maximal ideals. For example, if $n = 2$ then $U(\mathfrak{sl}_3^+)$ admits generators x, y, z , satisfying the relations:

$$xz = zx, \quad yz = zy, \quad xy - yx = z.$$

The center of $U(\mathfrak{sl}_3^+)$ is the polynomial algebra in the central variable z , and $U(\mathfrak{sl}_3^+)/(z-1)$ is isomorphic to the first Weyl algebra over the ground field. Here, the quantum scenario differs from the classical one: it is well-known that there

*Work supported by Fundação para a Ciência e a Tecnologia (FCT) through Centro de Matemática da Universidade do Porto.

are primitive ideals of $U_q(\mathfrak{sl}_3^+)$ which are not maximal (see [18], for example), and consequently $U_q(\mathfrak{g}^+)$ has non-simple primitive quotients, in general.

Let \mathbb{K} be an algebraically closed field of characteristic 0 and assume $q \in \mathbb{K}^*$ is not a root of unity. Then, $U_q(\mathfrak{g}^+)$ is the \mathbb{K} -algebra with generators e_1, \dots, e_n , which satisfy the so-called quantum Serre relations:

$$\begin{aligned} e_i e_j - e_j e_i &= 0 & \text{if } |i - j| \neq 1, \\ e_i^2 e_j - (q + q^{-1}) e_i e_j e_i + e_j e_i^2 &= 0 & \text{if } |i - j| = 1. \end{aligned}$$

The center of $U_q(\mathfrak{g}^+)$ was computed by Alev and Dumas [1], and by Caldero [5, 6]. It is the polynomial algebra over \mathbb{K} in the central indeterminates z_1, \dots, z_l , where $l = \lfloor \frac{n+1}{2} \rfloor$. If $n = 2$, then $U_q(\mathfrak{sl}_3^+)$ can be presented by generators X, Y, Z , satisfying the relations:

$$ZX = q^{-1}XZ, \quad ZY = qYZ, \quad XY - q^{-1}YX = Z.$$

The center of $U_q(\mathfrak{sl}_3^+)$ is the polynomial algebra in the variable $z_1 = (XY - qYX)Z$.

In [14], Kirkman and Small showed that $(z_1 - 1)$ is a maximal ideal of $U_q(\mathfrak{sl}_3^+)$ and that the factor algebra $A_q = U_q(\mathfrak{sl}_3^+)/ (z_1 - 1)$, in spite of not being isomorphic to the Weyl algebra $\mathbb{A}_1(\mathbb{K})$, shares a number of ring theoretical properties with it: it is a simple Noetherian domain with trivial center, Gelfand-Kirillov dimension 2 and Krull dimension 1. In the first part of this paper, we generalize these results of Kirkman and Small to $U_q(\mathfrak{g}^+)$, for any $n \geq 2$. Specifically, we show that $(z_1 - \alpha_1, \dots, z_l - \alpha_l)$ is a maximal ideal of $U_q(\mathfrak{g}^+)$ for any $\alpha_1, \dots, \alpha_l \in \mathbb{K}^*$, and conclude that the corresponding factor algebra is a simple Noetherian domain with trivial center and even Gelfand-Kirillov dimension, which is not isomorphic to a Weyl algebra over \mathbb{K} .

We then proceed to the second part of the paper, where we study the (left) primitive ideals of $U_q(\mathfrak{sl}_4^+)$ in full detail. To make use of the stratification theory of Goodearl and Letzter [10], we consider the natural action of the 3-torus $\mathcal{H} = (\mathbb{K}^*)^3$ on $U_q(\mathfrak{sl}_4^+)$. Relative to this action, the prime spectrum of $U_q(\mathfrak{sl}_4^+)$ is partitioned into $4! = 24$ strata (as shown, in a much more general context, in [12]), given in Proposition 4.1. By analyzing the maximal portion of each stratum, we obtain all primitive ideals of $U_q(\mathfrak{sl}_4^+)$, and we also compute their heights using the catenarity of $U_q(\mathfrak{sl}_4^+)$ and Tauvel's height formula (see [9]). This was achieved by Malliavin [18] for $U_q(\mathfrak{sl}_3^+)$ and, recently, by Launois [15] for $U_q(\mathfrak{so}_5^+)$. In the latter case, Launois succeeded in computing the automorphism group of $U_q(\mathfrak{so}_5^+)$, using partial results of Andruskiewitsch and Dumas [3]. A noteworthy by-product of our study is that the Gelfand-Kirillov dimension of the primitive factors of $U_q(\mathfrak{sl}_4^+)$ is one of 0, 2 or 4. In particular, as in the classical case, this dimension is always even.

Finally, we construct a simple $U_q(\mathfrak{sl}_4^+)$ -module with annihilator P , for any primitive ideal P of $U_q(\mathfrak{sl}_4^+)$. This, of course, doesn't exhaust the simple $U_q(\mathfrak{sl}_4^+)$ -modules, but it solves the problem of deciding whether or not an element belongs to a specified primitive ideal of $U_q(\mathfrak{sl}_4^+)$, and thus makes it quite easy to distinguish between the primitive ideals of this algebra.

Part of the results of this paper were obtained while we were at the University of Wisconsin-Madison, preparing our doctoral dissertation. We would like to express our gratitude to Georgia Benkart, for all her help and expert advising.

2 Basic set-up

We work over an algebraically closed field \mathbb{K} of characteristic 0 and fix a parameter $q \in \mathbb{K}^*$ which is not a root of unity. Let $\mathfrak{g} = \mathfrak{sl}_{n+1}$ be the complex semisimple Lie algebra of traceless $(n+1) \times (n+1)$ matrices and consider its maximal nilpotent subalgebra $\mathfrak{g}^+ = \mathfrak{sl}_{n+1}^+$ consisting of the strictly upper triangular matrices in \mathfrak{g} . As usual, $[k] = \frac{q^k - q^{-k}}{q - q^{-1}}$ is the q -version of the integer $k \in \mathbb{Z}$.

2.1 The algebra $U_q(\mathfrak{g}^+)$

The quantized enveloping algebra $U_q(\mathfrak{g}^+)$ is the associative unital \mathbb{K} -algebra given by the Chevalley generators e_1, \dots, e_n , subject to the quantum Serre relations

$$e_i e_j - e_j e_i = 0 \quad \text{if } |i - j| \neq 1, \quad (1)$$

$$e_i^2 e_j - (q + q^{-1}) e_i e_j e_i + e_j e_i^2 = 0 \quad \text{if } |i - j| = 1. \quad (2)$$

Let $Q = \mathbb{Z}^n$ be the free abelian group of rank n with canonical basis $\{\alpha_1, \dots, \alpha_n\}$, and $Q^+ = \mathbb{N}^n$ be its submonoid. There is a nondegenerate bilinear form on $Q \times Q$ determined by $(\alpha_i, \alpha_j) = 2, -1$ or 0 if $i = j, |i - j| = 1$ or $|i - j| > 1$, respectively. By the homogeneity of the quantum Serre relations it follows that $U_q(\mathfrak{g}^+)$ has a Q^+ -grading given by assigning degree α_i to the generator e_i . We use the terminology *weight* instead of degree for this grading and write $wt(u) = \beta$ if $u \in U_q(\mathfrak{g}^+)$ has weight $\beta \in Q^+$.

2.2 PBW basis

As in [20, App. 2], we recursively define weight elements $X_{ij}, 1 \leq i < j \leq n+1$, by setting $X_{i,i+1} = e_i$ and $X_{ij} = X_{ik} X_{kj} - q^{-1} X_{kj} X_{ik}$, for $1 \leq i < k < j \leq n+1$ (this is independent of the choice of k). Note that $wt(X_{ij}) = \alpha_i + \dots + \alpha_{j-1}$, for $i < j$. The set $\{X_{ij}\}$ can be linearly ordered using the rule

$$X_{ij} < X_{kl} \iff (k < i) \text{ or } (k = i \text{ and } l < j),$$

and we use the alternative notation X_k for the k th element in this increasing chain, so that $\{X_{ij}\}_{1 \leq i < j \leq n+1} = \{X_k\}_{1 \leq k \leq m}$, where $m = \frac{1}{2}n(n+1)$. We also write $X^{\mathbf{b}} = X_1^{b_1} \dots X_m^{b_m}$, for $\mathbf{b} = (b_1, \dots, b_m) \in \mathbb{N}^m$.

The following results of Ringel are well-known.

Theorem 2.1 ([20, Thm. 2, Cor.]).

- (a) *The algebra $U_q(\mathfrak{g}^+)$ is an iterated skew polynomial ring of the form*

$$\mathbb{K}[X_1][X_2; \tau_2, \delta_2] \cdots [X_m; \tau_m, \delta_m],$$

where τ_j is an algebra automorphism and δ_j is a \mathbb{K} -linear τ_j -derivation;

- (b) *The monomials in $\{X^{\mathbf{b}} \mid \mathbf{b} \in \mathbb{N}^m\}$ form a basis of $U_q(\mathfrak{g}^+)$, and for all $i < j$ we have*

$$X_j X_i = q^{v_{ji}} X_i X_j + r,$$

where $v_{ji} = (wt(X_i), wt(X_j))$ and r is a linear combination of monomials in X_{i+1}, \dots, X_{j-1} ;

- (c) *The prime ideals of $U_q(\mathfrak{g}^+)$ are completely prime.*

2.3 The degree of an element of $U_q(\mathfrak{g}^+)$

Define an order relation on \mathbb{N}^m by $\mathbf{b} < \mathbf{c} \iff$ there is $1 \leq k \leq m$ such that $b_k < c_k$ and $b_t = c_t$ for all $t > k$. Along with Theorem 2.1(b), this determines an increasing filtration $\{\mathcal{F}_{\mathbf{a}}\}_{\mathbf{a} \in \mathbb{N}^m}$ of $U_q(\mathfrak{g}^+)$ given by

$$\mathcal{F}_{\mathbf{a}} = \bigoplus_{\mathbf{b} \leq \mathbf{a}} \mathbb{K} \cdot X^{\mathbf{b}},$$

and the corresponding graded algebra is the quantum affine space with generators $\theta_1, \dots, \theta_m$ and relations $\theta_j \theta_i = q^{v_{ji}} \theta_i \theta_j$ for $i < j$, where $\theta_i = gr X_i$ and $v_{ji} = (wt(X_i), wt(X_j))$.

For $u \in U_q(\mathfrak{g}^+)$, set $deg(u) = \mathbf{a} \in \mathbb{N}^m$ if $u \neq 0$ and \mathbf{a} is the unique element of \mathbb{N}^m that satisfies $u \in \mathcal{F}_{\mathbf{a}}$ and $u \notin \mathcal{F}_{\mathbf{b}}$ for any $\mathbf{b} < \mathbf{a}$. We say that u has degree \mathbf{a} . Note that $deg(uv) = deg(u) + deg(v)$ for all nonzero $u, v \in U_q(\mathfrak{g}^+)$.

2.4 Normal elements of $U_q(\mathfrak{g}^+)$

According to work of Alev and Dumas [1], and Caldero [6, 7], there exist weight elements $\Delta_1, \dots, \Delta_n$ of $U_q(\mathfrak{g}^+)$ such that the following theorem holds.

Theorem 2.2 ([6, 7]). *For $1 \leq i, j \leq n$, we have:*

- (a) $e_i \Delta_j = q^{\delta_{ij} - \delta_{i, n+1-j}} \Delta_j e_i$;
- (b) *The subalgebra of $U_q(\mathfrak{g}^+)$ generated by the Δ_i is a (commutative) polynomial algebra $\mathbb{K}[\Delta_1, \dots, \Delta_n]$ in n variables;*
- (c) *The center $Z_q(\mathfrak{g}^+)$ of $U_q(\mathfrak{g}^+)$ is the polynomial algebra in the variables $\{\Delta_k \Delta_{n+1-k} \mid 1 \leq k \leq n/2\}$ if n is even and $\{\Delta_k \Delta_{n+1-k} \mid 1 \leq k \leq (n-1)/2\} \cup \{\Delta_{(n+1)/2}\}$ if n is odd.*

It was noted in [16, 4.4] that

$$gr(\Delta_i) = gr(X_{i, n+1}) gr(X_{i-1, n}) \cdots gr(X_{2, n+3-i}) gr(X_{1, n+2-i}), \quad (3)$$

in the graded algebra of 2.3.

We fix some more notation for the entire paper. The center of $U_q(\mathfrak{g}^+)$ is denoted by $Z_q(\mathfrak{g}^+)$ and $l = \lfloor \frac{n+1}{2} \rfloor$. The elements z_1, \dots, z_l are defined by

$$z_i = \begin{cases} \Delta_i \Delta_{n+1-i} & \text{if } i < l, \\ \Delta_l \Delta_{l+1} & \text{if } i = l \text{ and } n \text{ is even,} \\ \Delta_l & \text{if } i = l \text{ and } n \text{ is odd,} \end{cases}$$

so that $Z_q(\mathfrak{g}^+) = \mathbb{K}[z_1, \dots, z_l]$. The integer m is the number of positive roots of the Lie algebra \mathfrak{sl}_{n+1} , i.e., $m = \frac{1}{2}n(n+1)$.

2.5 The prime and primitive ideals of $U_q(\mathfrak{g}^+)$

In this brief paragraph we summarize a portion of stratification theory of Good-earl and Letzter that is essential for our study. The reader should refer to [10] for further details, all proofs, and an explanation of the terminology. Only left primitive ideals are considered; in view of the antiautomorphism $e_i \mapsto e_i$ of $U_q(\mathfrak{g}^+)$, this is not a serious restriction.

Let \mathcal{H} be the n -torus $(\mathbb{K}^*)^n$. Corresponding to each $\bar{\lambda} = (\lambda_1, \dots, \lambda_n) \in \mathcal{H}$ there is an automorphism $\sigma_{\bar{\lambda}}$ of $U_q(\mathfrak{g}^+)$ given by $\sigma_{\bar{\lambda}}(e_i) = \lambda_i e_i$, for all $1 \leq i \leq n$. This defines a rational action of \mathcal{H} on $U_q(\mathfrak{g}^+)$ such that the induced grading of $U_q(\mathfrak{g}^+)$ by the character group of \mathcal{H} coincides with the weight space decomposition of 2.1 (identifying the i th projection map $\bar{\lambda} \mapsto \lambda_i$ with the simple root α_i). Since \mathcal{H} acts by automorphisms, the action carries over to the spaces $\text{Spec } U_q(\mathfrak{g}^+)$ and $\text{Prim } U_q(\mathfrak{g}^+)$ of prime and primitive ideals of $U_q(\mathfrak{g}^+)$, respectively, equipped with the Jacobson topology. Let $\mathcal{H}\text{-Spec } U_q(\mathfrak{g}^+) \subseteq \text{Spec } U_q(\mathfrak{g}^+)$ be the subspace of \mathcal{H} -invariant prime ideals, that is, $\mathcal{H}\text{-Spec } U_q(\mathfrak{g}^+)$ consists of the prime ideals J of $U_q(\mathfrak{g}^+)$ that are generated by weight elements of $U_q(\mathfrak{g}^+)$. By [10, Prop. 4.2] and Theorem 2.1, $\mathcal{H}\text{-Spec } U_q(\mathfrak{g}^+)$ is a finite set consisting of the ideals of the form $(P : \mathcal{H}) := \bigcap_{h \in \mathcal{H}} h.P$, for $P \in \text{Spec } U_q(\mathfrak{g}^+)$.

The above determines a decomposition of $\text{Spec } U_q(\mathfrak{g}^+)$ into \mathcal{H} -strata:

$$\text{Spec } U_q(\mathfrak{g}^+) = \bigcup_{J \in \mathcal{H}\text{-Spec } U_q(\mathfrak{g}^+)} \text{Spec}_J U_q(\mathfrak{g}^+),$$

where $\text{Spec}_J U_q(\mathfrak{g}^+) = \{P \in \text{Spec } U_q(\mathfrak{g}^+) \mid (P : \mathcal{H}) = J\}$ is the \mathcal{H} -stratum of J in $\text{Spec } U_q(\mathfrak{g}^+)$, and similarly for $\text{Prim } U_q(\mathfrak{g}^+)$. It follows from [10, Thm. 4.4] that the primitive ideals of $U_q(\mathfrak{g}^+)$ are the maximal elements of $\text{Spec}_J U_q(\mathfrak{g}^+)$, for each $J \in \mathcal{H}\text{-Spec } U_q(\mathfrak{g}^+)$. Furthermore, since \mathbb{K} is algebraically closed, [10, Thm. 2.6 or Thm. 6.8] imply that \mathcal{H} acts transitively on each of the sets $\text{Prim}_J U_q(\mathfrak{g}^+)$. Therefore, the \mathcal{H} -orbits of primitive ideals of $U_q(\mathfrak{g}^+)$ are parametrized by the elements of the set $\mathcal{H}\text{-Spec } U_q(\mathfrak{g}^+)$, which has cardinality $(n+1)!$ by work of Gorelik [12, Prop. 5.3.3] (see also [3, 3.4.1]).

Given $J \in \mathcal{H}\text{-Spec } U_q(\mathfrak{g}^+)$, let Ξ_J be the set of nonzero weight elements of $U_q(\mathfrak{g}^+)/J$, with respect to the Q^+ -grading inherited from $U_q(\mathfrak{g}^+)$. The following result of Goodearl and Letzter describes the \mathcal{H} -strata of $U_q(\mathfrak{g}^+)$.

Theorem 2.3 ([10, Thm. 6.6]). *Let J , Ξ_J and $\text{Spec}_J U_q(\mathfrak{g}^+)$ be as above. Then Ξ_J is an Ore set in $U_q(\mathfrak{g}^+)/J$. If $U_q(\mathfrak{g}^+)_J$ denotes the localization of $U_q(\mathfrak{g}^+)/J$ at Ξ_J , we have:*

- (a) *The localization map $U_q(\mathfrak{g}^+) \rightarrow U_q(\mathfrak{g}^+)/J \rightarrow U_q(\mathfrak{g}^+)_J$ induces a homeomorphism of $\text{Spec}_J U_q(\mathfrak{g}^+)$ onto $\text{Spec } U_q(\mathfrak{g}^+)_J$.*
- (b) *Contraction and extension induce mutually inverse homeomorphisms between $\text{Spec } U_q(\mathfrak{g}^+)_J$ and $\text{Spec } Z(U_q(\mathfrak{g}^+)_J)$, where $Z(U_q(\mathfrak{g}^+)_J)$ is the center of $U_q(\mathfrak{g}^+)_J$.*

3 Generalizations of the Weyl-Hayashi algebra

If $n = 2$ then $Z_q(\mathfrak{sl}_3^+)$ is a polynomial algebra in the central variable z_1 . The factor algebra $U_q(\mathfrak{sl}_3^+)/(z_1 - 1)$ was introduced by Hayashi in [13], in connection with oscillator representations of quantized enveloping algebras of semisimple Lie algebras of types A and C . In [14], Kirkman and Small showed that $U_q(\mathfrak{sl}_3^+)/(z_1 - 1)$ is a simple Noetherian domain of Gelfand-Kirillov dimension 2, which is not isomorphic to the Weyl algebra $\mathbb{A}_1(\mathbb{K})$ (see also [2] and [18]). The relevance of their result is that the primitive factor algebras of the enveloping algebra of a finite-dimensional nilpotent Lie algebra over a field of characteristic 0 are isomorphic to Weyl algebras over the base field (see [8, Thm. 4.7.9]). In

particular, those primitive factor algebras of Gelfand-Kirillov dimension 2 must be isomorphic to $\mathbb{A}_1(\mathbb{K})$. In this section we generalize the result of Kirkman and Small to $U_q(\mathfrak{g}^+)$, thus proposing other analogues of the Weyl algebras $\mathbb{A}_k(\mathbb{K})$.

3.1 Gröbner bases

Let us introduce some basic techniques from the theory of Gröbner bases. We follow [4]. Recall the filtration, associated graded algebra and the notion of degree defined in 2.3 in terms of the PBW basis of $U_q(\mathfrak{g}^+)$. Given a subset F of $U_q(\mathfrak{g}^+)$, set $\deg(F) = \{\deg(f) \mid 0 \neq f \in F\} \subseteq \mathbb{N}^m$. It is clear that if L is a left, right, or two-sided ideal of $U_q(\mathfrak{g}^+)$, then $\deg(L)$ is stable under translation by elements of \mathbb{N}^m ; in other words, $\deg(L)$ is a *monoideal* of \mathbb{N}^m . The set $\{f_1, \dots, f_s\} \subseteq L$ is said to be a *Gröbner basis* for L if (see [4, Def. 2.8])

$$\deg(L) = \bigcup_{j=1}^s (\deg(f_j) + \mathbb{N}^m).$$

Recall also from 2.4 that the center of $U_q(\mathfrak{g}^+)$ is $Z_q(\mathfrak{g}^+) = \mathbb{K}[z_1, \dots, z_l]$. Let $\mathbf{t} = (t_1, \dots, t_l) \in \mathbb{K}^l$ and set $f_j^{\mathbf{t}} = z_j - t_j$, $I^{\mathbf{t}} = \sum_{j=1}^l U_q(\mathfrak{g}^+)f_j^{\mathbf{t}}$ and $\tilde{I}^{\mathbf{t}} = \sum_{j=1}^l Z_q(\mathfrak{g}^+)f_j^{\mathbf{t}}$.

Proposition 3.1. *The set $\{f_1^{\mathbf{t}}, \dots, f_l^{\mathbf{t}}\}$ is a Gröbner basis for $I^{\mathbf{t}}$.*

Proof. It is enough to show that

$$\deg(I^{\mathbf{t}}) \subseteq \bigcup_{j=1}^l (\deg(f_j^{\mathbf{t}}) + \mathbb{N}^m),$$

since the other inclusion follows from the fact that $\deg(I^{\mathbf{t}})$ is a monoideal and $\{f_1^{\mathbf{t}}, \dots, f_l^{\mathbf{t}}\} \subseteq I^{\mathbf{t}}$.

Recall that by our separation of variables results [16, Lem. 1, Thm. 2], there is a set \mathcal{M} consisting of monomials $X^{\mathbf{a}}$ ($\mathbf{a} \in \mathbb{N}^m$) in the PBW basis elements X_1, \dots, X_m such that

$$U_q(\mathfrak{g}^+) = \bigoplus_{u \in \mathcal{M}} uZ_q(\mathfrak{g}^+).$$

Furthermore, by construction,

$$X^{\mathbf{a}} \in \mathcal{M} \implies \mathbf{a} - \deg(z_j) \notin \mathbb{N}^m, \quad \text{for all } 1 \leq j \leq l. \quad (4)$$

Thus,

$$I^{\mathbf{t}} = U_q(\mathfrak{g}^+)\tilde{I}^{\mathbf{t}} = \bigoplus_{u \in \mathcal{M}} u\tilde{I}^{\mathbf{t}}. \quad (5)$$

Assume $u, u' \in \mathcal{M}$, $p, p' \in \tilde{I}^{\mathbf{t}}$ and $\deg(up) = \deg(u'p')$. By (3), the elements $\deg(z_j)$, $1 \leq j \leq l$ are free generators of the monoid $\deg(Z_q(\mathfrak{g}^+))$, and in particular $\deg(\tilde{I}^{\mathbf{t}}) \subseteq \bigoplus_{i=1}^l \mathbb{N}\deg(z_i)$. Let $\mathbf{a}, \mathbf{b} \in \mathbb{N}^m$ be such that $u = X^{\mathbf{a}}$ and $u' = X^{\mathbf{b}}$. There exist $n_i, n'_i \in \mathbb{N}$, $1 \leq i \leq l$, with

$$\mathbf{a} + \sum_{i=1}^l n_i \deg(z_i) = \mathbf{b} + \sum_{i=1}^l n'_i \deg(z_i).$$

Assume $\mathbf{a} \neq \mathbf{b}$. Then there is $1 \leq i \leq l$ such that $n_i \neq n'_i$. Without loss of generality assume $n_1 < n'_1$. Then,

$$\mathbf{a} - \deg(z_1) + \sum_{i=2}^l n_i \deg(z_i) = \mathbf{b} + n'_1 \deg(z_1) + \sum_{i=2}^l n'_i \deg(z_i), \quad (6)$$

for $n'_1 = (n'_1 - n_1 - 1) \in \mathbb{N}$. But (6) implies that $\mathbf{a} - \deg(z_1) \in \mathbb{N}^m$ by (3), and this contradicts (4). Thus $\mathbf{a} = \mathbf{b}$ and $u = u'$.

Let $0 \neq f \in I^{\mathbf{t}}$. Then by (5) we can write $f = \sum_{j=1}^k u_j p_j$ with $u_j \in \mathcal{M}$, $0 \neq p_j \in \tilde{I}^{\mathbf{t}}$ and the u_j pairwise distinct, $1 \leq j \leq k$. The discussion above shows that $\deg(f) = \max_j \deg(u_j p_j)$, with respect to the total order on \mathbb{N}^m defined in 2.3. Say $\deg(f) = \deg(u_1 p_1)$. We have

$$\deg(p_1) \in \deg(\tilde{I}^{\mathbf{t}}) \subseteq \left(\bigoplus_{i=1}^l \mathbb{N} \deg(z_i) \right) \setminus \{\bar{0}\} \subseteq \bigcup_{i=1}^l (\deg(z_i) + \mathbb{N}^m),$$

since $\tilde{I}^{\mathbf{t}}$ is a proper ideal of $Z_q(\mathfrak{g}^+)$. Finally, as $\deg(f_i^{\mathbf{t}}) = \deg(z_i)$ for all $1 \leq i \leq l$, we obtain the desired conclusion:

$$\deg(f) = \deg(u_1) + \deg(p_1) \in \bigcup_{i=1}^l (\deg(f_i^{\mathbf{t}}) + \mathbb{N}^m).$$

□

For $0 \neq f \in U_q(\mathfrak{g}^+)$ write $f = \sum_{\mathbf{a} \in \mathbb{N}^m} c_{\mathbf{a}} X^{\mathbf{a}}$, where $c_{\mathbf{a}} \in \mathbb{K}$ and the sum is finite. Define $\mathcal{N}(f) = \{\mathbf{a} \in \mathbb{N}^m \mid c_{\mathbf{a}} \neq 0\}$. We are going to use the division algorithm developed in [4, Thm. 2.1] to prove the next result.

Corollary 3.2. *For all $\mathbf{t} \in \mathbb{K}^l$, the ideal $I^{\mathbf{t}}$ of $U_q(\mathfrak{g}^+)$ is semiprime.*

Proof. Using for example [11, Thm. 2.7], it is enough to show $f^2 \in I^{\mathbf{t}} \implies f \in I^{\mathbf{t}}$, for all $f \in U_q(\mathfrak{g}^+)$. By the (left) division algorithm of [4, Thm. 2.1], there exist elements $g_1, \dots, g_l, r \in U_q(\mathfrak{g}^+)$ (unique under certain conditions) such that

$$f = \sum_{i=1}^l g_i f_i^{\mathbf{t}} + r,$$

with either $r = 0$ or $\mathcal{N}(r) \subseteq \mathbb{N}^m \setminus \bigcup_{i=1}^l (\deg(f_i^{\mathbf{t}}) + \mathbb{N}^m)$. If $r = 0$ there is nothing to prove, so assume $r \neq 0$. Since

$$I^{\mathbf{t}} \ni f^2 = \sum_{i,j=1}^l g_i f_i^{\mathbf{t}} g_j f_j^{\mathbf{t}} + \sum_{j=1}^l r g_j f_j^{\mathbf{t}} + \sum_{i=1}^l g_i f_i^{\mathbf{t}} r + r^2$$

and the $f_i^{\mathbf{t}}$ are central, it must be that $r^2 \in I^{\mathbf{t}}$. If $\deg(r) = (a_1, \dots, a_m)$, then $\deg(r^2) = (2a_1, \dots, 2a_m) \in \deg(I^{\mathbf{t}})$. By Proposition 3.1, there are $1 \leq i \leq l$ and $\mathbf{b} \in \mathbb{N}^m$ such that $\deg(r^2) = \deg(f_i^{\mathbf{t}}) + \mathbf{b}$. But by (3), $\deg(f_i^{\mathbf{t}})$ is a string of zeros and ones, so it follows that $\deg(r) \in \deg(f_i^{\mathbf{t}}) + \mathbb{N}^m$, which contradicts the assumption on $\mathcal{N}(r)$. Therefore $r = 0$ and $f \in I^{\mathbf{t}}$. □

3.2 The \mathcal{H} -stratum of (0)

We want to describe the space of all primitive ideals of $U_q(\mathfrak{sl}_{n+1})^+$ that do not contain nonzero weight elements, i.e. $\text{Prim}_{(0)} U_q(\mathfrak{g}^+)$. By [10, Thm. 4.4] and Theorem 2.3, this space is homeomorphic to the space of maximal ideals of $Z(\mathcal{Q})$, where \mathcal{Q} is the localization of $U_q(\mathfrak{g}^+)$ at the Ore set $\Xi_{(0)}$ of nonzero weight elements of $U_q(\mathfrak{g}^+)$, and $Z(\mathcal{Q})$ is the center of \mathcal{Q} .

Lemma 3.3. *The center of \mathcal{Q} is the commutative Laurent polynomial algebra*

$$Z(\mathcal{Q}) = \mathbb{K}[z_1^{\pm 1}, \dots, z_l^{\pm 1}].$$

Proof. Let Λ be the set of nonzero weight elements of $Z_q(\mathfrak{g}^+)$. First we want to show that $Z(\mathcal{Q}) = Z_q(\mathfrak{g}^+)[\Lambda^{-1}]$, the localization of the center of $U_q(\mathfrak{g}^+)$ at Λ . Fix $b^{-1}a \in Z(\mathcal{Q})$, where $a, b \in U_q(\mathfrak{g}^+)$ with b a nonzero weight element. Without loss of generality, it can be assumed that a is a weight element, as $Z(\mathcal{Q})$ is graded by Q . Consider the set $L = \{u \in U_q(\mathfrak{g}^+) \mid ub^{-1}a \in U_q(\mathfrak{g}^+)\}$. As $b^{-1}a$ is central, L is a (nonzero) two-sided ideal of $U_q(\mathfrak{g}^+)$. We can use [6, Thm. 5.2] to conclude that there is a nonzero $x \in Z_q(\mathfrak{g}^+)$ such that $xb^{-1}a \in U_q(\mathfrak{g}^+)$. Once more, by weight considerations, x can be assumed to be a weight element, so that $x \in \Lambda$. Therefore, $b^{-1}a = x^{-1}(xb^{-1}a)$ with $x \in \Lambda$ and $xb^{-1}a \in Z_q(\mathfrak{g}^+)$. The above shows that $Z(\mathcal{Q}) \subseteq Z_q(\mathfrak{g}^+)[\Lambda^{-1}]$. The reverse inclusion is obvious.

Now we determine the set Λ . It is easy to see, for example from [6, 3.1], that $wt(z_1), \dots, wt(z_l)$ are \mathbb{Z} -independent elements of Q^+ ; in fact, $wt(z_j) = 2(\varpi_j + \varpi_{n+1-j})$ for $1 \leq j \leq l$, where $\varpi_1, \dots, \varpi_n$ are the fundamental weights of \mathfrak{g} . Since $Z_q(\mathfrak{g}^+)$ is the polynomial algebra $\mathbb{K}[z_1, \dots, z_l]$, it follows that the weight elements of $Z_q(\mathfrak{g}^+)$ are precisely the scalar multiples of the monomials $z_1^{\alpha_1} \cdots z_l^{\alpha_l}$. Thus, $Z_q(\mathfrak{g}^+)[\Lambda^{-1}] = \mathbb{K}[z_1^{\pm 1}, \dots, z_l^{\pm 1}]$, and the lemma is proved. \square

By the above lemma, the maximal ideals of $Z(\mathcal{Q})$ are those of the form

$$\sum_{j=1}^l Z(\mathcal{Q})(z_j - t_j), \quad \text{for } \mathbf{t} = (t_1, \dots, t_l) \in (\mathbb{K}^*)^l. \quad (7)$$

Fix $\mathbf{t} \in (\mathbb{K}^*)^l$. The ideal (7) corresponds to the maximal ideal $\sum_{j=1}^l \mathcal{Q}(z_j - t_j)$ of \mathcal{Q} , by extension (see Theorem 2.3(b)), and the latter corresponds to the ideal

$$T^{\mathbf{t}} := \left(\sum_{j=1}^l \mathcal{Q}(z_j - t_j) \right) \cap U_q(\mathfrak{g}^+) \quad (8)$$

of $U_q(\mathfrak{g}^+)$ under the homeomorphism of Theorem 2.3(a) with $J = (0)$. Thus $T^{\mathbf{t}}$ is maximal within the \mathcal{H} -stratum of (0) in $\text{Spec } U_q(\mathfrak{g}^+)$, and by [10, Thm. 4.4], $T^{\mathbf{t}}$ is a primitive ideal of $U_q(\mathfrak{g}^+)$ containing no nonzero weight elements. Furthermore, by construction, any primitive ideal of $U_q(\mathfrak{g}^+)$ with this property is of the form $T^{\mathbf{t}'}$, for some $\mathbf{t}' \in (\mathbb{K}^*)^l$.

Recall the definition of $I^{\mathbf{t}}$ in 3.1. The next proposition gives the ideal $T^{\mathbf{t}}$ a more familiar form.

Proposition 3.4. *Let $\mathbf{t} \in (\mathbb{K}^*)^l$. Then the ideal $I^{\mathbf{t}}$ is primitive. In fact,*

$$T^{\mathbf{t}} = I^{\mathbf{t}} = \sum_{j=1}^l U_q(\mathfrak{g}^+)(z_j - t_j).$$

Proof. Assume $\mathbf{t} \in (\mathbb{K}^*)^l$. Note that the inclusion $I^{\mathbf{t}} \subseteq T^{\mathbf{t}}$ is clear. Let P be any prime ideal of $U_q(\mathfrak{g}^+)$ containing $I^{\mathbf{t}}$, and suppose P contains a nonzero weight element. Then $(P : \mathcal{H}) \neq (0)$, and hence there is a nonzero weight element $g \in (P : \mathcal{H}) \cap Z_q(\mathfrak{g}^+)$, by [6, Thm. 5.2] and the fact that $(P : \mathcal{H})$ is a graded ideal. In particular, $g \in P$. We have

$$P \cap Z_q(\mathfrak{g}^+) \supseteq I^{\mathbf{t}} \cap Z_q(\mathfrak{g}^+),$$

and $g \in P \cap Z_q(\mathfrak{g}^+) \setminus I^{\mathbf{t}} \cap Z_q(\mathfrak{g}^+)$, as $I^{\mathbf{t}}$ contains no nonzero weight elements since $I^{\mathbf{t}} \subseteq T^{\mathbf{t}}$. But

$$I^{\mathbf{t}} \cap Z_q(\mathfrak{g}^+) = \sum_{j=1}^l Z_q(\mathfrak{g}^+)(z_j - t_j)$$

is a maximal ideal of $Z_q(\mathfrak{g}^+)$, and therefore $P \cap Z_q(\mathfrak{g}^+) = Z_q(\mathfrak{g}^+)$. The latter is a contradiction since $1 \notin P$. This shows that $(P : \mathcal{H}) = (0)$ and thus $P \in \text{Spec}_{(0)} U_q(\mathfrak{g}^+)$. Hence P extends to a prime ideal P^e of \mathcal{Q} with

$$P^e \supseteq \sum_{j=1}^l \mathcal{Q}(z_j - t_j), \quad (9)$$

as $P \supseteq I^{\mathbf{t}}$. The ideal on the right-hand side of (9) is a maximal ideal of \mathcal{Q} , and thus equality must hold in (9). Therefore, P is the contraction to $U_q(\mathfrak{g}^+)$ of $\sum_{j=1}^l \mathcal{Q}(z_j - t_j)$, which by definition is $T^{\mathbf{t}}$.

We have seen that $T^{\mathbf{t}}$ is the unique prime ideal of $U_q(\mathfrak{g}^+)$ containing $I^{\mathbf{t}}$. However, it was shown in Corollary 3.2 that $I^{\mathbf{t}}$ is an intersection of prime ideals, which forces $I^{\mathbf{t}} = T^{\mathbf{t}}$. \square

We remark that it is necessary to require that $\mathbf{t} \in (\mathbb{K}^*)^l$ for Proposition 3.4 to hold. If, for example, $t_1 = 0$ and $n \geq 2$, then $I^{\mathbf{t}}$ is not even a prime ideal (even though it remains semiprime by Corollary 3.2), since $\Delta_1 \Delta_n = z_1 \in I^{\mathbf{t}}$ and yet neither Δ_1 nor Δ_n is in $I^{\mathbf{t}}$, by Proposition 3.1. The only possible exception to this type of counterexample would be to have $t_l = 0$ and n odd, as in this case $z_l = \Delta_l$; we have not investigated this situation here.

Theorem 3.5. *Assume $\mathbf{t} \in (\mathbb{K}^*)^l$. Then $I^{\mathbf{t}}$ is a maximal ideal of $U_q(\mathfrak{g}^+)$. It is also minimal among primitive ideals of $U_q(\mathfrak{g}^+)$.*

Proof. Assume $I^{\mathbf{t}}$ is not maximal. Then there is a maximal ideal P of $U_q(\mathfrak{g}^+)$ such that $I^{\mathbf{t}} \subsetneq P \subsetneq U_q(\mathfrak{g}^+)$. Since $(I^{\mathbf{t}} : \mathcal{H}) = (0)$ and $I^{\mathbf{t}}$ is primitive by Proposition 3.4, it follows that $I^{\mathbf{t}}$ is a maximal element of $\text{Spec}_{(0)} U_q(\mathfrak{g}^+)$, by [10, Thm. 4.4]. Hence $(P : \mathcal{H}) \neq (0)$. As in the proof of Proposition 3.4, this yields a contradiction. Thus $I^{\mathbf{t}}$ is indeed a maximal ideal.

The other statement of the theorem follows because $U_q(\mathfrak{g}^+)$ is an iterated skew polynomial ring over \mathbb{K} , hence a constructible \mathbb{K} -algebra (see [19, 9.4.12]), and by [19, Thm. 9.4.21] it satisfies the Nullstellensatz over the algebraically closed field \mathbb{K} . \square

3.3 Gelfand-Kirillov dimension of $U_q(\mathfrak{g}^+)/I^{\mathfrak{t}}$

It is a simple matter to compute the GK (Gelfand-Kirillov) dimension of the factor algebra $U_q(\mathfrak{g}^+)/I^{\mathfrak{t}}$. We can use Proposition 3.1 along with the techniques of [4, Sec. 4] for example, as we did in [17, 5.2.3]. Another approach is via [9, Thm. 4.8], where the authors show that $U_q(\mathfrak{g}^+)$ is catenary and that Tauvel's height formula holds in $U_q(\mathfrak{g}^+)$. By the latter property, given $P \in \text{Spec } U_q(\mathfrak{g}^+)$,

$$\text{GKdim}(U_q(\mathfrak{g}^+)/P) = \text{GKdim}(U_q(\mathfrak{g}^+)) - \text{height}(P) = m - \text{height}(P). \quad (10)$$

If $\mathfrak{t} \in (\mathbb{K}^*)^l$ and $J \subseteq I^{\mathfrak{t}}$ is a prime ideal, then clearly $J \in \text{Spec}_{(0)} U_q(\mathfrak{g}^+)$. Since the spaces $\text{Spec}_{(0)} U_q(\mathfrak{g}^+)$ and $\text{Spec } Z(\mathcal{Q})$ are homeomorphic, it follows that the height of $I^{\mathfrak{t}}$ equals the height of the corresponding maximal ideal (7) of $Z(\mathcal{Q})$, which is l by Lemma 3.3. Therefore, (10) yields $\text{GKdim}(U_q(\mathfrak{g}^+)/I^{\mathfrak{t}}) = m - l$ for $\mathfrak{t} \in (\mathbb{K}^*)^l$, a result which holds more generally for all $\mathfrak{t} \in \mathbb{K}^l$, as seen in [17, 5.2.3].

Corollary 3.6. *Let $\mathfrak{t} \in (\mathbb{K}^*)^l$. The factor algebra $U_q(\mathfrak{g}^+)/I^{\mathfrak{t}}$ is a simple Noetherian domain with center \mathbb{K} and GK dimension $m - l$. In particular, the GK dimension of $U_q(\mathfrak{g}^+)/I^{\mathfrak{t}}$ is always even but $U_q(\mathfrak{g}^+)/I^{\mathfrak{t}}$ is not isomorphic to a Weyl algebra $A_k(\mathbb{K})$ for any $k \geq 1$.*

Proof. The statement about the center of $U_q(\mathfrak{g}^+)/I^{\mathfrak{t}}$ is a consequence of the Nullstellensatz for $U_q(\mathfrak{g}^+)$ (see the proof of Theorem 3.5 for more details). The last statement follows since Δ_1 is invertible in $U_q(\mathfrak{g}^+)/I^{\mathfrak{t}}$, as $\Delta_1 \Delta_n - t_1 \in I^{\mathfrak{t}}$ and $t_1 \neq 0$, whereas the only invertible elements of $A_k(\mathbb{K})$ are the nonzero scalars. \square

4 The primitive ideals of $U_q(\mathfrak{sl}_4^+)$

In this section we study the primitive spectrum of $U_q(\mathfrak{sl}_4^+)$ in full detail. This was done for $U_q(\mathfrak{sl}_3^+)$ in [18], and for $U_q(\mathfrak{so}_5^+)$ in [15]. We begin by determining the set $\mathcal{H}\text{-Spec } U_q(\mathfrak{sl}_4^+)$ of all \mathcal{H} -invariant (completely) prime ideals of $U_q(\mathfrak{sl}_4^+)$. It is a finite set of cardinality $4! = 24$ whose elements parametrize the \mathcal{H} -orbits of primitive ideals of $U_q(\mathfrak{sl}_4^+)$, by the stratification theory of Goodearl and Letzter [10] and work of Gorelik [12]. Then, we explicitly describe each \mathcal{H} -stratum in $\text{Prim } U_q(\mathfrak{sl}_4^+)$, compute the height of all primitive ideals of $U_q(\mathfrak{sl}_4^+)$, and give an example of a simple $U_q(\mathfrak{sl}_4^+)$ -module with annihilator P , for each $P \in \text{Prim } U_q(\mathfrak{sl}_4^+)$. An interesting outcome of this analysis is that all primitive factor algebras of $U_q(\mathfrak{sl}_4^+)$ have even GK dimension, as occurs in the classical case.

4.1 Structure of $U_q(\mathfrak{sl}_4^+)$

A PBW basis of $U_q(\mathfrak{sl}_4^+)$, as seen in 2.2, is given by

$$\begin{aligned} X_1 &= e_3, & X_2 &= e_2 e_3 - q^{-1} e_3 e_2, & X_3 &= e_2, \\ X_4 &= e_1 X_2 - q^{-1} X_2 e_1, & X_5 &= e_1 e_2 - q^{-1} e_2 e_1, & X_6 &= e_1, \end{aligned}$$

and we can take

$$\Delta_1 = X_4, \quad \Delta_2 = X_2 X_5 - q^{-1} X_3 X_4,$$

$$\Delta_3 = q^{-2} \left((q - q^{-1})^2 X_1 X_3 X_6 - (q - q^{-1}) X_1 X_5 - (q - q^{-1}) X_2 X_6 + X_4 \right),$$

$$z_1 = \Delta_1 \Delta_3, \quad z_2 = \Delta_2,$$

so that $Z_q(\mathfrak{sl}_4^+) = \mathbb{K}[z_1, z_2]$.

Consider the *diagram automorphism* η of $U_q(\mathfrak{sl}_4^+)$, where $\eta(e_i) = e_{4-i}$ for $i = 1, 2, 3$. The element Δ_3 above was defined so that $\eta(\Delta_1) = \Delta_3$.

4.2 \mathcal{H} -Spec $U_q(\mathfrak{sl}_4^+)$

In [17, 5.3.3] we used the ideas of Goodearl and Letzter to obtain Proposition 4.1 below; specifically, we used [10, Lem. 3.2] and the proofs of [10, Lem. 3.3, Prop. 3.4]. As it is known that $|\mathcal{H}\text{-Spec } U_q(\mathfrak{sl}_4^+)| = 4!$, another approach to proving this result would be to show that the ideals listed therein are distinct prime ideals of $U_q(\mathfrak{sl}_4^+)$, as they are clearly \mathcal{H} -invariant. Notice that the automorphism η defined in 4.1 acts on $\mathcal{H}\text{-Spec } U_q(\mathfrak{sl}_4^+)$ and each η -orbit has size either one or two, since $\eta^2 = 1$.

Proposition 4.1. *The space $\mathcal{H}\text{-Spec } U_q(\mathfrak{sl}_4^+)$ consists of the 24 ideals in the following list, where ideals that are in the same η -orbit have been grouped together:*

1. (0) ;
2. (Δ_2) ;
3. $(\Delta_1), (\Delta_3)$;
4. (Δ_1, Δ_3) ;
5. $(e_1 e_2 - q e_2 e_1), (e_2 e_3 - q^{-1} e_3 e_2)$;
6. $(e_1 e_2 - q^{-1} e_2 e_1), (e_2 e_3 - q e_3 e_2)$;
7. $(e_1), (e_3)$;
8. $(e_1 e_2 - q e_2 e_1, e_2 e_3 - q e_3 e_2), (e_1 e_2 - q^{-1} e_2 e_1, e_2 e_3 - q^{-1} e_3 e_2)$;
9. $(e_1 e_2 - q^{-1} e_2 e_1, e_2 e_3 - q e_3 e_2)$;
10. $(e_1 e_2 - q e_2 e_1, e_2 e_3 - q^{-1} e_3 e_2)$;
11. $(e_2 e_3 - q e_3 e_2, e_1), (e_1 e_2 - q^{-1} e_2 e_1, e_3)$;
12. $(e_2 e_3 - q^{-1} e_3 e_2, e_1), (e_1 e_2 - q e_2 e_1, e_3)$;
13. $(e_1, e_2), (e_2, e_3)$;
14. (e_2) ;
15. (e_1, e_3) ;
16. (e_1, e_2, e_3) .

4.3 Prim $U_q(\mathfrak{sl}_4^+)$

Finally, we can determine all primitive ideals of $U_q(\mathfrak{sl}_4^+)$ and compute their heights by studying each of the spaces $\text{Prim}_J U_q(\mathfrak{sl}_4^+)$, for all possible choices of $J \in \mathcal{H}\text{-Spec } U_q(\mathfrak{sl}_4^+)$. We denote the ideals given by Proposition 4.1 by J_i , $J_{i,a}$ or $J_{i,b}$ according to their position in that list, so that $J_{13,a} = (e_1, e_2)$, $J_{13,b} = (e_2, e_3)$ and $J_{14} = (e_2)$, for example. If M is a $U_q(\mathfrak{sl}_4^+)$ -module, we denote its annihilator by $\text{ann } M$.

Recall from [1] that the center of $U_q(\mathfrak{sl}_3^+)$ is the polynomial algebra in the quantum Casimir element

$$\Omega = (\dot{e}_1 \dot{e}_2 - q^{-1} \dot{e}_2 \dot{e}_1)(\dot{e}_1 \dot{e}_2 - q \dot{e}_2 \dot{e}_1) \in U_q(\mathfrak{sl}_3^+), \quad (11)$$

where \dot{e}_1 and \dot{e}_2 denote the Chevalley generators of $U_q(\mathfrak{sl}_3^+)$. For $i = 1, 2$ let S_i be the subalgebra of $U_q(\mathfrak{sl}_4^+)$ generated by e_i and e_{i+1} . Then $S_i \simeq U_q(\mathfrak{sl}_3^+)$ and therefore the center of S_i is the polynomial algebra $\mathbb{K}[\Omega_i]$ in the variable

$$\Omega_i = (e_i e_{i+1} - q^{-1} e_{i+1} e_i)(e_i e_{i+1} - q e_{i+1} e_i). \quad (12)$$

4.3.1 Annihilators of the finite-dimensional simple $U_q(\mathfrak{sl}_4^+)$ -modules: the \mathcal{H} -strata of (e_2) , (e_1, e_2) , (e_2, e_3) , (e_1, e_3) and (e_1, e_2, e_3)

As a result of all prime ideals of $U_q(\mathfrak{sl}_4^+)$ being completely prime, and given the assumption that \mathbb{K} is algebraically closed, it can be shown that the simple finite-dimensional $U_q(\mathfrak{sl}_4^+)$ -modules are one-dimensional. If $V = \mathbb{K}v_0$ is such a one-dimensional module, then there exist scalars $\alpha_i \in \mathbb{K}$ with either $\alpha_2 = 0$, or $\alpha_2 \neq 0$ and $\alpha_1 = 0 = \alpha_3$, satisfying $e_i.v_0 = \alpha_i v_0$, for $i = 1, 2, 3$ (the conditions on the scalars α_i follow directly from the quantum Serre relations (2)). Consequently, the primitive ideals that occur as annihilators of finite-dimensional simple $U_q(\mathfrak{sl}_4^+)$ -modules are the maximal ideals of the form $(e_1 - \alpha_1, e_2 - \alpha_2, e_3 - \alpha_3)$, with the α_i as above; these belong to one of the following \mathcal{H} -strata: (e_1, e_2, e_3) , (e_1, e_2) , (e_2, e_3) , (e_1, e_3) , (e_2) .

Proposition 4.2. (a) $\text{Prim}_{(e_1, e_2, e_3)} U_q(\mathfrak{sl}_4^+) = \{(e_1, e_2, e_3)\}$;

(b) $\text{Prim}_{(e_1, e_2)} U_q(\mathfrak{sl}_4^+) = \{(e_1, e_2, e_3 - \alpha) \mid \alpha \in \mathbb{K}^*\}$;

(c) $\text{Prim}_{(e_2, e_3)} U_q(\mathfrak{sl}_4^+) = \{(e_1 - \alpha, e_2, e_3) \mid \alpha \in \mathbb{K}^*\}$;

(d) $\text{Prim}_{(e_1, e_3)} U_q(\mathfrak{sl}_4^+) = \{(e_1, e_2 - \alpha, e_3) \mid \alpha \in \mathbb{K}^*\}$;

(e) $\text{Prim}_{(e_2)} U_q(\mathfrak{sl}_4^+) = \{(e_1 - \alpha, e_2, e_3 - \beta) \mid \alpha, \beta \in \mathbb{K}^*\}$.

The primitive ideals described above have height 6.

Proof. Parts (a)–(e) follow easily from the discussion above and the Nullstellensatz. The last statement follows from Tauvel’s height formula (10) since $\text{GKdim}(U_q(\mathfrak{sl}_4^+)) = 6$ and $\text{GKdim}(U_q(\mathfrak{sl}_4^+)/P) = 0$ for any of the primitive ideals listed in (a)–(e). \square

4.3.2 The \mathcal{H} -stratum of (0)

Proposition 4.3. *Let z_1 and z_2 be the generators of the center of $U_q(\mathfrak{sl}_4^+)$, as defined in 4.1. Then,*

$$\text{Prim}_{(0)} U_q(\mathfrak{sl}_4^+) = \{(z_1 - \alpha, z_2 - \beta) \mid \alpha, \beta \in \mathbb{K}^*\} \quad (13)$$

and these primitive ideals have height 2.

Proof. The equality (13) was proved in Proposition 3.4, and the height of the ideal $(z_1 - \alpha, z_2 - \beta)$, for $\alpha, \beta \in \mathbb{K}^*$, was computed in 3.3. \square

In [16, 6.3] we defined $U_q(\mathfrak{sl}_4^+)$ -modules $M_{(\alpha, \beta)}$, for $(\alpha, \beta) \in \mathbb{K}^2$, which were shown to be pairwise non-isomorphic, and simple under the assumption that $\alpha \neq 0$. It was also shown that $(z_1 - \alpha^2, z_2 - \beta) \subseteq \text{ann } M_{(\alpha, \beta)}$. In particular, if $\alpha, \beta \in \mathbb{K}^*$ and α' is a square root of α in \mathbb{K} , then

$$\text{ann } M_{(\alpha', \beta)} = (z_1 - \alpha, z_2 - \beta), \quad (14)$$

since the ideal on the right-hand side of (14) is maximal.

4.3.3 The \mathcal{H} -stratum of (Δ_2)

Let $Q \in \text{Prim}_{(\Delta_2)} U_q(\mathfrak{sl}_4^+)$. Since $U_q(\mathfrak{sl}_4^+)$ satisfies the Nullstellensatz over \mathbb{K} (see the proof of Theorem 3.5), there exist scalars $\alpha, \beta \in \mathbb{K}$ such that $z_1 - \alpha, z_2 - \beta \in Q$; hence $\beta = 0$ as, by hypothesis, $z_2 = \Delta_2 \in Q$. If $\alpha = 0$ then $z_1 = \Delta_1 \Delta_3 \in Q$, which implies that $\Delta_1 \in (\Delta_2)$ or $\Delta_3 \in (\Delta_2)$ because z_1 is a weight element and Q is completely prime. This is a contradiction, as $(\Delta_i, \Delta_2) \neq (\Delta_2)$ for any $i = 1, 3$, and hence $z_1 - \alpha \in Q$ for some $\alpha \in \mathbb{K}^*$. Indeed, let $U_q(\mathfrak{sl}_4^+)$ act on the vector space $A = \mathbb{K}[x, y]$ of all polynomials in the variables x and y , by the formulas ($a, b \geq 0$):

$$\begin{aligned} e_1.x^a y^b &= [a]x^{a-1}y^b, \\ e_2.x^a y^b &= x^{a+1}y^{b+1}, \\ e_3.x^a y^b &= [b]x^a y^{b-1}, \end{aligned} \quad (15)$$

where $e_1.y^b = 0 = e_3.x^a$.

Lemma 4.4. *The formulas (15) above endow A with the structure of a simple $U_q(\mathfrak{sl}_4^+)$ -module with $\text{ann } A \in \text{Prim}_{(\Delta_2)} U_q(\mathfrak{sl}_4^+)$. We also have $z_1 - q^{-2} \in \text{ann } A$.*

Proof. It is easy to check that the formulas in (15) extend to give an action of $U_q(\mathfrak{sl}_4^+)$ on A , and that A thus becomes a simple $U_q(\mathfrak{sl}_4^+)$ -module. Also, straightforward computations yield:

$$\begin{aligned} \Delta_1.x^a y^b &= -q^{a-b-1}x^a y^b, \\ \Delta_2.x^a y^b &= 0, \\ \Delta_3.x^a y^b &= -q^{b-a-1}x^a y^b, \end{aligned} \quad \text{for all } a, b \geq 0.$$

Therefore, $z_1 - q^{-2}, z_2 \in \text{ann } A$. Let $P = \text{ann } A$. We use Proposition 4.1 to see that $P \in \text{Prim}_{(\Delta_2)} U_q(\mathfrak{sl}_4^+)$: noting that

$$P \cap \{e_1, e_2, e_3, e_1 e_2 - q^{\pm 1} e_2 e_1, e_2 e_3 - q^{\pm 1} e_3 e_2, \Delta_1, \Delta_3\} = \emptyset$$

and $\Delta_2 \in P$ (recall that $P = \text{ann } A$ and it is thus easy to check if a given element is in P), it must be that $(P : \mathcal{H}) = (\Delta_2)$. \square

Proposition 4.5. *Let $P = \text{ann } A$, as before. Then,*

$$\text{Prim}_{(\Delta_2)} U_q(\mathfrak{sl}_4^+) = \{\sigma_{\bar{\lambda}}(P) \mid \bar{\lambda} \in (\mathbb{K}^*)^3\} \quad (16)$$

and these primitive ideals have height 2.

Proof. We have already seen that $P \in \text{Prim}_{(\Delta_2)} U_q(\mathfrak{sl}_4^+)$ in Lemma 4.4. Since $\mathcal{H} = (\mathbb{K}^*)^3$ acts transitively on $\text{Prim}_{(\Delta_2)} U_q(\mathfrak{sl}_4^+)$, the first statement follows. To finish the proof, it is enough to show that the primitive ideal P has height 2, as \mathcal{H} acts by automorphisms. Note that $z_1 - q^{-2} \in P \setminus (\Delta_2)$ and so the chain

$$(0) \subsetneq (\Delta_2) \subsetneq P$$

gives $\text{height}(P) \geq 2$. By Tauvel's height formula (10),

$$\text{height}(P) = 6 - \text{GKdim}(U_q(\mathfrak{sl}_4^+)/P)$$

and it remains to show that $\text{GKdim}(U_q(\mathfrak{sl}_4^+)/P) \geq 4$. We do so by exhibiting a subalgebra of $U_q(\mathfrak{sl}_4^+)/P$ with GK dimension 4.

Let $S = S_2$ (as defined in 4.3) and T be the subalgebra of $U_q(\mathfrak{sl}_4^+)$ generated by e_2, e_3 and $\Delta_1 = X_4$. Then T is the Ore extension $S[\Delta_1; \tau]$, where τ is the automorphism of S determined by $\tau(e_2) = e_2$ and $\tau(e_3) = qe_3$; in particular, $\text{GKdim}(T) = 4$. Consider the inclusion $\iota : S \rightarrow T$, and the map $\phi : T \rightarrow U_q(\mathfrak{sl}_4^+)/P$ obtained by composing the inclusion $T \rightarrow U_q(\mathfrak{sl}_4^+)$ and the natural epimorphism $U_q(\mathfrak{sl}_4^+) \rightarrow U_q(\mathfrak{sl}_4^+)/P$. Since P is a completely prime ideal of $U_q(\mathfrak{sl}_4^+)$, it follows that $\ker \phi = P \cap T$ and $\ker \phi \circ \iota = P \cap S$ are (completely) prime ideals of T and S , respectively. In [18, Thm. 2.4] the lattice of prime ideals of $U_q(\mathfrak{sl}_3^+)$ was obtained, and we can use that result to conclude, via the natural isomorphism $S \simeq U_q(\mathfrak{sl}_3^+)$, that $P \cap S = (0)$, as

$$\{e_2e_3 - q^{\pm 1}e_3e_2, \Omega_2 - \alpha \mid \alpha \in \mathbb{K}\} \cap P = \emptyset.$$

Let $P' = P \cap T$ and assume $P' \neq (0)$. Take

$$p = u_0 + u_1\Delta_1 + \cdots + u_k\Delta_1^k \in P' \setminus (0)$$

with $k \geq 0$ minimal such that $u_0, \dots, u_k \in S$ and $u_k \neq 0$. Since Ω_2 is central in S and $\Omega_2\Delta_1 = q^{-2}\Delta_1\Omega_2$, we have

$$\left[(1 - q^{-2k})u_0 + \cdots + (q^{-2(k-1)} - q^{-2k})u_{k-1}\Delta_1^{k-1} \right] \Omega_2 = \Omega_2 p - q^{-2k}p\Omega_2 \in P'.$$

As P' is completely prime and $\Omega_2 \notin P$, the minimality of k implies that $u_r = 0$ for all $r < k$. Hence $p = u_k\Delta_1^k$ and again because $\Delta_1 \notin P$, it must be that $k = 0$ and $p = u_0$. Thus $p \in P \cap S = (0)$, a contradiction. This proves that ϕ is injective and, as a consequence,

$$\text{GKdim}(U_q(\mathfrak{sl}_4^+)/P) \geq \text{GKdim}(T) = 4.$$

□

The task of finding a simple $U_q(\mathfrak{sl}_4^+)$ -module with given annihilator $Q \in \text{Prim}_{(\Delta_2)} U_q(\mathfrak{sl}_4^+)$ is now trivial: if $Q = \sigma_{\bar{\lambda}}(P)$, then Q is the annihilator of the simple module obtained by twisting A by the automorphism $\sigma_{\bar{\lambda}}^{-1} = \sigma_{\bar{\lambda}^{-1}}$.

4.3.4 The \mathcal{H} -strata of (Δ_1) and (Δ_3)

Consider the $U_q(\mathfrak{sl}_4^+)$ -module $P_{(\alpha,\beta,\gamma)} = \mathbb{K}[x, y^{\pm 1}]$, defined in [16, 6.3] in terms of the parameters $\alpha, \beta, \gamma \in \mathbb{K}$. Taking $\alpha = 0, \beta = 1$ and $\gamma = q^{-1}$, we obtain the following formulas for the action of $U_q(\mathfrak{sl}_4^+)$ on $P_{(0,1,q^{-1})}$:

$$\begin{aligned} e_1.x^a y^b &= \begin{cases} q^{-(a+b)}(x^a y^b + [a]x^{a-1}y^{b-1}) & \text{if } b \geq 1 \\ q^{-(a+b)}x^a y^b + [a]x^{a-1}y^{b-1} & \text{if } b \leq 0, \end{cases} \\ e_2.x^a y^b &= \begin{cases} q^b x^{a+1} y^b & \text{if } b \geq 0 \\ x^{a+1} y^b & \text{if } b \leq 0, \end{cases} \\ e_3.x^a y^b &= \begin{cases} -q^{a-b}[a]x^{a-1}y^{b+1} & \text{if } b \geq 0 \\ -[a]x^{a-1}y^{b+1} & \text{if } b \leq -1. \end{cases} \end{aligned}$$

Lemma 4.6. *The module $P_{(0,1,q^{-1})}$ is simple and $P \in \text{Prim}_{(\Delta_1)} U_q(\mathfrak{sl}_4^+)$, where $P = \text{ann } P_{(0,1,q^{-1})}$.*

Proof. First, we show that $P_{(0,1,q^{-1})}$ is simple. Let $(0) \neq W \subseteq P_{(0,1,q^{-1})}$ be a submodule and take $p = p(x, y) \in W \setminus \{0\}$ such that the x -degree of p is as small as possible, say $a \geq 0$. If $a > 0$, then $e_3.p$ is a nonzero element of W with smaller x -degree, which is a contradiction. Thus $a = 0$ and $p = p(y) \in \mathbb{K}[y^{\pm 1}]$. Notice that $e_3 e_2.y^k \in \mathbb{K}^*.y^{k+1}$ for all $k \in \mathbb{Z}$ and hence, acting by a high enough power of $e_3 e_2$, we can assume that $p \in \mathbb{K}[y]$ and choose such a nonzero element of W with minimum y -degree, say $b \geq 0$. Assume $b \geq 1$ and write

$$p = c_0 + \cdots + c_{b-1}y^{b-1} + y^b$$

with all $c_i \in \mathbb{K}$. Since $e_1.y^k = q^{-k}y^k$ for all $k \in \mathbb{Z}$, we have

$$c_{b-1}(q^{-b} - q^{-(b-1)})y^{b-1} + \cdots + c_0(q^{-b} - 1) = q^{-b}p - e_1.p \in W. \quad (17)$$

Given the minimality of b , the element in (17) must be equal to 0; hence $c_r = 0$ for all $0 \leq r < b$ and $p = y^b$. Computing still, we obtain

$$y^{b-1} = (qe_1 e_2 - q^{-b}e_2).y^b \in W,$$

which contradicts the minimality of b . Therefore $b = 0$ and $1 \in W$. By construction, the modules $P_{(\alpha,\beta,\gamma)}$ are generated by $1 \in \mathbb{K}[x, y^{\pm 1}]$ and hence $W = P_{(0,1,q^{-1})}$, proving the simplicity of $P_{(0,1,q^{-1})}$.

Let $P = \text{ann } P_{(0,1,q^{-1})}$. We have just seen that P is a primitive ideal of $U_q(\mathfrak{sl}_4^+)$; thus $P \in \text{Prim}_J U_q(\mathfrak{sl}_4^+)$, for some $J \in \mathcal{H}\text{-Spec } U_q(\mathfrak{sl}_4^+)$. It is easy to check that none of the following weight elements is in P : $e_1, e_2, e_3, e_1 e_2 - q^{\pm 1}e_2 e_1, e_2 e_3 - q^{\pm 1}e_3 e_2$ (neither, except for e_3 , annihilate $1 \in P_{(0,1,q^{-1})}$, and e_3 does not annihilate x , for example). Therefore, by Proposition 4.1, J must be one of the following ideals $(0), (\Delta_1), (\Delta_2), (\Delta_3)$ or (Δ_1, Δ_3) . Again by [16, 6.3] (or by direct computation) we see that $\Delta_1 \in P$ and $\Delta_2 - q^{-1} \in P$, and we can also check that $\Delta_3 \notin P$ (for example, $\Delta_3.1 = -(q - q^{-1})y$). Thus $J = (\Delta_1)$. \square

Let $Q_{(0,1,q^{-1})}$ be the $U_q(\mathfrak{sl}_4^+)$ -module obtained by twisting $P_{(0,1,q^{-1})}$ by the diagram automorphism η of 4.1. Recall that if $P_{(0,1,q^{-1})}$ is given by the representation ρ , then $Q_{(0,1,q^{-1})}$ is defined by the representation $\rho \circ \eta$. The following lemma is an easy consequence of Lemma 4.6.

Lemma 4.7. *The module $Q_{(0,1,q^{-1})}$ is simple and $Q \in \text{Prim}_{(\Delta_3)} U_q(\mathfrak{sl}_4^+)$, where $Q = \text{ann } Q_{(0,1,q^{-1})}$.*

Proof. The simplicity of $Q_{(0,1,q^{-1})}$ follows from Lemma 4.6 and the fact that η is onto. In particular, Q is a primitive ideal and $Q = \ker \rho \circ \eta = \eta(P)$, as $\eta^2 = 1$. Furthermore,

$$\sigma_{(\lambda_1, \lambda_2, \lambda_3)} \circ \eta = \eta \circ \sigma_{(\lambda_3, \lambda_2, \lambda_1)}, \quad \text{for all } (\lambda_1, \lambda_2, \lambda_3) \in \mathcal{H}, \quad (18)$$

and thus $(Q : \mathcal{H}) = \eta((P : \mathcal{H})) = (\Delta_3)$, since $\eta(\Delta_1) = \Delta_3$. Therefore, $Q \in \text{Prim}_{(\Delta_3)} U_q(\mathfrak{sl}_4^+)$. \square

Proposition 4.8. *Let $P = \text{ann } P_{(0,1,q^{-1})}$ and $Q = \text{ann } Q_{(0,1,q^{-1})}$. Then*

$$(a) \text{Prim}_{(\Delta_1)} U_q(\mathfrak{sl}_4^+) = \{\sigma_{\bar{\lambda}}(P) \mid \bar{\lambda} \in (\mathbb{K}^*)^3\},$$

$$(b) \text{Prim}_{(\Delta_3)} U_q(\mathfrak{sl}_4^+) = \{\sigma_{\bar{\lambda}}(Q) \mid \bar{\lambda} \in (\mathbb{K}^*)^3\},$$

and these primitive ideals have height 2.

Proof. Parts (a) and (b) follow from the previous two lemmas and the transitivity of the action of $\mathcal{H} = (\mathbb{K}^*)^3$ on each \mathcal{H} -stratum in $\text{Prim } U_q(\mathfrak{sl}_4^+)$. To prove the last statement it is enough to show that $\text{height}(Q) = 2$. The chain of prime ideals

$$(0) \subsetneq (\Delta_3) \subsetneq Q$$

is proper, as $\Delta_2 - q^{-1} = \eta(\Delta_2 - q^{-1}) \in Q \setminus (\Delta_3)$, so $\text{height}(Q) \geq 2$. The proof that $\text{height}(Q) \leq 2$ is done exactly as in Proposition 4.5, by using Tauvel's height formula and showing that $U_q(\mathfrak{sl}_4^+)/Q$ has a subalgebra of GK dimension 4. The same algebras S and T used in the proof of that proposition work in this setting, so we do not repeat the argument. \square

4.3.5 The \mathcal{H} -stratum of (Δ_1, Δ_3)

Let $B = \mathbb{K}[t]$ be the vector space of all polynomials in the variable t . It is easy to see that the formulas

$$\begin{aligned} e_1.t^k &= [k]t^{k-1}, \\ e_2.t^k &= t^{k+1}, \\ e_3.t^k &= [k]t^{k-1}, \end{aligned}$$

define an action of $U_q(\mathfrak{sl}_4^+)$ on B . Let $P = \text{ann } B$.

Lemma 4.9. *With the action described above, B becomes a simple $U_q(\mathfrak{sl}_4^+)$ -module and $P \in \text{Prim}_{(\Delta_1, \Delta_3)} U_q(\mathfrak{sl}_4^+)$.*

Proof. It is routine to check that B is simple; therefore the ideal P is primitive and $P \in \text{Prim}_J U_q(\mathfrak{sl}_4^+)$, for some $J \in \mathcal{H}\text{-Spec } U_q(\mathfrak{sl}_4^+)$. As none of the weight elements $e_1, e_2, e_3, e_1e_2 - q^{\pm 1}e_2e_1, e_2e_3 - q^{\pm 1}e_3e_2, \Delta_2$ annihilates B , yet Δ_1 and Δ_3 do annihilate it, it must be that $J = (\Delta_1, \Delta_3)$, as desired. \square

Recall the quantum Casimir element $\Omega \in U_q(\mathfrak{sl}_3^+)$ and the elements $\Omega_i, i = 1, 2$, of $U_q(\mathfrak{sl}_4^+)$ defined in 4.3. Let $P' = (\Delta_1, \Delta_3, e_1 - e_3, \Omega_1 - 1)$. By computing, we check that

$$e_1 - e_3, \Omega_1 - 1 \in P = \text{ann } B,$$

and consequently $P' \subseteq P$.

Lemma 4.10. *Let P' be as before. Then,*

$$U_q(\mathfrak{sl}_3^+)/(\Omega - 1) \simeq U_q(\mathfrak{sl}_4^+)/P', \quad (19)$$

and $P = P'$.

Proof. The inclusion $U_q(\mathfrak{sl}_3^+) \rightarrow U_q(\mathfrak{sl}_4^+)$, $e_i \mapsto e_i$, $i = 1, 2$, induces an algebra homomorphism $\phi : U_q(\mathfrak{sl}_3^+) \rightarrow U_q(\mathfrak{sl}_4^+)/P'$, which is onto because $e_1 - e_3 \in P'$. Moreover, since $\phi(\Omega - 1) = (\Omega_1 - 1) + P' = P'$, it follows that $\Omega - 1 \in \ker \phi$. By Theorem 3.5, $(\Omega - 1)$ is a maximal ideal of $U_q(\mathfrak{sl}_3^+)$; thus $\ker \phi = (\Omega - 1)$ and ϕ induces the desired isomorphism (19). In particular, $U_q(\mathfrak{sl}_4^+)/P'$ is a simple algebra and hence P' is a maximal ideal of $U_q(\mathfrak{sl}_4^+)$. This shows that $P' = P$, as we had already observed that $P' \subseteq P$. \square

Proposition 4.11. *The \mathcal{H} -stratum of (Δ_1, Δ_3) in $\text{Prim } U_q(\mathfrak{sl}_4^+)$ is*

$$\text{Prim}_{(\Delta_1, \Delta_3)} U_q(\mathfrak{sl}_4^+) = \{(\Delta_1, \Delta_3, e_1 - \alpha e_3, \Omega_1 - \beta) \mid \alpha, \beta \in \mathbb{K}^*\},$$

and these primitive ideals have height 4.

Proof. We have seen that $P = (\Delta_1, \Delta_3, e_1 - e_3, \Omega_1 - 1) \in \text{Prim}_{(\Delta_1, \Delta_3)} U_q(\mathfrak{sl}_4^+)$. Since \mathbb{K} is algebraically closed, the space $\text{Prim}_{(\Delta_1, \Delta_3)} U_q(\mathfrak{sl}_4^+)$ consists of a single \mathcal{H} -orbit and it is easy to check that

$$\mathcal{H}.P = \{(\Delta_1, \Delta_3, e_1 - \alpha e_3, \Omega_1 - \beta) \mid \alpha, \beta \in \mathbb{K}^*\}.$$

The statement about the height of the primitive ideals in the \mathcal{H} -orbit of P follows from Tauvel's height formula and Lemma 4.10, as $\text{GKdim}(U_q(\mathfrak{sl}_3^+)/(\Omega - 1)) = 2$ by Corollary 3.6. \square

4.3.6 The \mathcal{H} -strata of $(e_1 e_2 - q^{\pm 1} e_2 e_1)$ and $(e_2 e_3 - q^{\pm 1} e_3 e_2)$

We consider only the \mathcal{H} -stratum of $J_{5,b} = (e_2 e_3 - q^{-1} e_3 e_2)$ in detail, the cases of the \mathcal{H} -strata of $J_{5,a}$, $J_{6,a}$ and $J_{6,b}$ being similar. The factor algebra $U_q(\mathfrak{sl}_4^+)/J_{5,b}$ is isomorphic to $R := S_1[Y; \nu]$, where ν is the algebra automorphism of S_1 given by $\nu(e_1) = e_1$ and $\nu(e_2) = q e_2$. Indeed, there is a surjective algebra homomorphism $\phi : U_q(\mathfrak{sl}_4^+) \rightarrow R$ such that $\phi(e_i) = e_i$, $i = 1, 2$, and $\phi(e_3) = Y$. As $e_2 e_3 - q^{-1} e_3 e_2 \in \ker \phi$, ϕ induces a surjective map, which we still denote by ϕ , $U_q(\mathfrak{sl}_4^+)/J_{5,b} \rightarrow R$. The natural map $S_1 \rightarrow U_q(\mathfrak{sl}_4^+) \rightarrow U_q(\mathfrak{sl}_4^+)/J_{5,b}$ can be extended to an algebra homomorphism $\psi : R \rightarrow U_q(\mathfrak{sl}_4^+)/J_{5,b}$ such that $\psi(Y) = e_3 + J_{5,b}$, by the universal property of Ore extensions. The maps ϕ and ψ just defined are inverses of each other; in particular, $\phi : U_q(\mathfrak{sl}_4^+)/J_{5,b} \rightarrow R$ is an algebra isomorphism.

The algebra R is Q^+ -graded so that ϕ becomes an isomorphism of graded algebras. Hence, the spaces $\text{Spec}_{J_{5,b}} U_q(\mathfrak{sl}_4^+)$ and $\text{Spec}_{(0)} R$ can be identified via ϕ .

Lemma 4.12. *The ideal $J_{5,b}$ of $U_q(\mathfrak{sl}_4^+)$ is primitive.*

Proof. It suffices to show that $\text{Spec}_{(0)} R = \{(0)\}$, as $\text{Prim}_{J_{5,b}} U_q(\mathfrak{sl}_4^+)$ consists of the maximal elements of $\text{Spec}_{J_{5,b}} U_q(\mathfrak{sl}_4^+)$, by the stratification theory of Goodearl and Letzter [10].

Let $P \in \text{Spec}_{(0)} R$ and assume $P \neq (0)$. As P is a completely prime ideal of R (because R is a homomorphic image of $U_q(\mathfrak{sl}_4^+)$ and the latter algebra has the property that all of its prime ideals are completely prime), it follows that $P' = P \cap S_1$ is a (completely) prime ideal of $S_1 \simeq U_q(\mathfrak{sl}_3^+)$. Since P contains no nonzero weight elements, the same is true of P' and by [18, Thm. 2.4], either $\Omega_1 - \alpha \in P'$ for some $\alpha \in \mathbb{K}^*$ or $P' = (0)$.

Assume $\Omega_1 - \alpha \in P'$, where $\alpha \in \mathbb{K}^*$. In R , we have

$$Y\Omega_1 = q^2\Omega_1Y, \quad (20)$$

and thus

$$(q^2 - 1)\alpha Y = Y(\Omega_1 - \alpha) - q^2(\Omega_1 - \alpha)Y \in P. \quad (21)$$

This is a contradiction, as Y is a nonzero weight element of R and (21) implies that $Y \in P$. Therefore $P' = (0)$.

Let $f \in P \setminus (0)$, say

$$f = u_0 + \cdots + u_k Y^k,$$

with $k \geq 0$, $u_0, \dots, u_k \in S_1$ and $u_k \neq 0$, and assume such an element was chosen with minimum k . Then, by (20) and the fact that Ω_1 is central in S_1 ,

$$[(1 - q^{2k})u_0 + \cdots + (1 - q^2)u_{k-1}Y^{k-1}] \Omega_1 = f\Omega_1 - q^{2k}\Omega_1 f \in P. \quad (22)$$

Since P is completely prime and $\Omega_1 \notin P$, the minimality of k implies that $u_r = 0$ for all $r < k$. Hence $f = u_k Y^k$ and again it must be that $u_k \in P$, as $Y \notin P$. Thus $u_k \in P \cap S_1 = (0)$, a contradiction. The contradiction resulted from our assumption that $P \neq (0)$, so $\text{Spec}_{(0)} R = \{(0)\}$ and $\text{Spec}_{J_{5,b}} U_q(\mathfrak{sl}_4^+) = \{J_{5,b}\}$. \square

Proposition 4.13. (a) $\text{Prim}_{J_{5,a}} U_q(\mathfrak{sl}_4^+) = \{(e_1 e_2 - q e_2 e_1)\}$;

(b) $\text{Prim}_{J_{5,b}} U_q(\mathfrak{sl}_4^+) = \{(e_2 e_3 - q^{-1} e_3 e_2)\}$;

(c) $\text{Prim}_{J_{6,a}} U_q(\mathfrak{sl}_4^+) = \{(e_1 e_2 - q^{-1} e_2 e_1)\}$;

(d) $\text{Prim}_{J_{6,b}} U_q(\mathfrak{sl}_4^+) = \{(e_2 e_3 - q e_3 e_2)\}$.

The primitive ideals described above have height 2.

Proof. Part (b) of the proposition is Lemma 4.12 and the others are similar (for example, for (a) we need only use the automorphism η). The height of these primitive ideals can be computed using Tauvel's height formula. For example, $\text{height}(J_{5,b}) = 2$ because $\text{GKdim}(U_q(\mathfrak{sl}_4^+)/J_{5,b}) = \text{GKdim}(R) = 4$. \square

To finish this paragraph, we give an example of a simple $U_q(\mathfrak{sl}_4^+)$ -module C with annihilator $J_{5,b}$. Simple $U_q(\mathfrak{sl}_4^+)$ -modules with annihilators $J_{5,a}$, $J_{6,a}$ and $J_{6,b}$ can be readily obtained by twisting C by η and/or replacing q by q^{-1} in the formulas below. Let $C = \mathbb{K}[x, y^{\pm 1}]$ be the vector space with basis $\{x^a y^b \mid a \geq 0, b \in \mathbb{Z}\}$, and define an action of $U_q(\mathfrak{sl}_4^+)$ on C by

$$\begin{aligned} e_1.x^a y^b &= [a]x^{a-1}y^b, \\ e_2.x^a y^b &= q^{-b}x^{a+1}y^{b-1}, \\ e_3.x^a y^b &= x^a y^{b+1}, \end{aligned} \quad a \geq 0, b \in \mathbb{Z}. \quad (23)$$

(We leave it as an exercise to show that (23) does give a well-defined action of $U_q(\mathfrak{sl}_4^+)$ on C .)

Lemma 4.14. *The $U_q(\mathfrak{sl}_4^+)$ -module C defined above is simple and $\text{ann } C = J_{5,b}$.*

Proof. Let $P = \text{ann } C$. The simplicity of C is easy to check, as it is to verify that $J_{5,b} \subseteq P$. Thus, $P \in \text{Prim}_J U_q(\mathfrak{sl}_4^+)$ for some $J_{5,b} \subseteq J \in \mathcal{H}\text{-Spec } U_q(\mathfrak{sl}_4^+)$. Since

$$\{e_1, e_2, e_3, e_1e_2 - q^{\pm 1}e_2e_1, e_2e_3 - qe_3e_2\} \cap P = \emptyset \quad (24)$$

and

$$e_2e_3 - q^{-1}e_3e_2 \notin (\Delta_1) \cup (\Delta_2) \cup (\Delta_3) \cup (\Delta_1, \Delta_3) \cup (0), \quad (25)$$

(note that (25) was verified when we studied each of the individual strata involved), it can only be that $J = J_{5,b}$, by Proposition 4.1, and therefore $P = J_{5,b}$, by Proposition 4.13. \square

4.3.7 The \mathcal{H} -strata of (e_1) and (e_3)

Since $U_q(\mathfrak{sl}_4^+)/\langle e_1 \rangle \simeq U_q(\mathfrak{sl}_3^+)$, the spaces $\text{Prim}_{(e_1)} U_q(\mathfrak{sl}_4^+)$ and $\text{Prim}_{(0)} U_q(\mathfrak{sl}_3^+)$ can be naturally identified. As we have seen in 3.2, or otherwise by [18, Thm. 2.4],

$$\text{Prim}_{(0)} U_q(\mathfrak{sl}_3^+) = \{(\Omega - \alpha) \mid \alpha \in \mathbb{K}^*\}, \quad (26)$$

where Ω is given by (11).

Proposition 4.15. *Let Ω_i , $i = 1, 2$, be defined as in (12). Then,*

$$(a) \text{Prim}_{(e_1)} U_q(\mathfrak{sl}_4^+) = \{(e_1, \Omega_2 - \alpha) \mid \alpha \in \mathbb{K}^*\};$$

$$(b) \text{Prim}_{(e_3)} U_q(\mathfrak{sl}_4^+) = \{(e_3, \Omega_1 - \alpha) \mid \alpha \in \mathbb{K}^*\}.$$

The primitive ideals described above have height 4.

Proof. Part (a) follows from (26). Since

$$U_q(\mathfrak{sl}_4^+)/\langle e_1, \Omega_2 - \alpha \rangle \simeq U_q(\mathfrak{sl}_3^+)/(\Omega - \alpha)$$

and $\text{GKdim}(U_q(\mathfrak{sl}_3^+)/(\Omega - \alpha)) = 2$, by Corollary 3.6, the last statement follows from Tauvel's height formula. Part (b) is analogous and can be obtained from (a) via the automorphism η . \square

An example of a simple $U_q(\mathfrak{sl}_3^+)$ -module with annihilator $(\Omega - 1) \subseteq U_q(\mathfrak{sl}_3^+)$ is the vector space $D = \mathbb{K}[t]$, with action induced by

$$\begin{aligned} e_1.t^k &= [k]t^{k-1}, \\ e_2.t^k &= t^{k+1}, \end{aligned} \quad k \geq 0$$

(see [16, 6.2]). This action extends to $U_q(\mathfrak{sl}_4^+)$ by defining $e_3.t^k = 0$ for all $k \geq 0$, and hence we obtain a simple $U_q(\mathfrak{sl}_4^+)$ -module with annihilator $(e_3, \Omega_1 - 1)$. Twisting this action by the automorphisms of the form $\sigma_{\bar{\lambda}}$, $\bar{\lambda} \in (\mathbb{K}^*)^3$, and η , we easily get examples of simple $U_q(\mathfrak{sl}_4^+)$ -modules corresponding to each of the primitive ideals of Proposition 4.15.

4.3.8 The \mathcal{H} -strata of $(e_1e_2 - q^{\pm 1}e_2e_1, e_2e_3 - q^{\pm 1}e_3e_2)$

Let $\delta, \epsilon \in \{-1, 1\}$, and consider the quantum affine space $\mathbb{K}_{\delta, \epsilon}[x, y, z]$, generated by x, y, z with relations $xz = zx, xy = q^\epsilon yx, yz = q^\delta zy$. There are isomorphisms

$$U_q(\mathfrak{sl}_4^+)/J_{8,a} \simeq \mathbb{K}_{1,1}[x, y, z], \quad U_q(\mathfrak{sl}_4^+)/J_{8,b} \simeq \mathbb{K}_{-1,-1}[x, y, z], \quad (27)$$

$$U_q(\mathfrak{sl}_4^+)/J_9 \simeq \mathbb{K}_{1,-1}[x, y, z], \quad U_q(\mathfrak{sl}_4^+)/J_{10} \simeq \mathbb{K}_{-1,1}[x, y, z], \quad (28)$$

each sending e_1, e_2, e_3 to x, y, z , respectively.

Denote the localization of $\mathbb{K}_{\delta, \epsilon}[x, y, z]$ at the multiplicatively closed set generated by the normal elements x, y and z by $\mathbb{K}_{\delta, \epsilon}[x^{\pm 1}, y^{\pm 1}, z^{\pm 1}]$. It follows from [10, Thm. 4.4] and Theorem 2.3 that the isomorphism $U_q(\mathfrak{sl}_4^+)/J_9 \simeq \mathbb{K}_{1,-1}[x, y, z]$ provides a homeomorphism between $\text{Prim}_{J_9} U_q(\mathfrak{sl}_4^+)$ and the space of maximal ideals of $\mathbb{K}_{1,-1}[x^{\pm 1}, y^{\pm 1}, z^{\pm 1}]$, and similarly for $J_{8,a}, J_{8,b}$ and J_{10} . Furthermore, by Theorem 2.3(b), contraction and extension induce mutually inverse homeomorphisms between the space of maximal ideals of $\mathbb{K}_{\delta, \epsilon}[x^{\pm 1}, y^{\pm 1}, z^{\pm 1}]$ and the space of maximal ideals of $Z_{\delta, \epsilon}$, the center of $\mathbb{K}_{\delta, \epsilon}[x^{\pm 1}, y^{\pm 1}, z^{\pm 1}]$.

Lemma 4.16. *The center of $\mathbb{K}_{\delta, \epsilon}[x^{\pm 1}, y^{\pm 1}, z^{\pm 1}]$ is a Laurent polynomial algebra in the variable $w_{\delta, \epsilon} = x^\delta z^\epsilon$:*

$$Z_{\delta, \epsilon} = \mathbb{K}[w_{\delta, \epsilon}^{\pm 1}].$$

Proof. Let $w \in \mathbb{K}_{\delta, \epsilon}[x^{\pm 1}, y^{\pm 1}, z^{\pm 1}]$ be nonzero. Then $w \in Z_{\delta, \epsilon}$ exactly when $dw = wd$ for all $d \in \{x, y, z\}$. Write

$$w = \sum_{(a,b,c) \in \mathbb{Z}^3} \lambda(a, b, c) x^a y^b z^c,$$

where the $\lambda(a, b, c) \in \mathbb{K}$ are all but finitely many equal to 0. A simple computation shows that $xw = wx$ is equivalent to the equation

$$\sum_{(a,b,c) \in \mathbb{Z}^3} \lambda(a, b, c) x^{a+1} y^b z^c = \sum_{(a,b,c) \in \mathbb{Z}^3} \lambda(a, b, c) q^{-\epsilon b} x^{a+1} y^b z^c, \quad (29)$$

and thus, since q is not a root of unity, $b = 0$ whenever $\lambda(a, b, c) \neq 0$. So, we can write

$$w = \sum_{(a,c) \in \mathbb{Z}^2} \lambda(a, c) x^a z^c,$$

and it is clear that w now commutes with both x and z . Similarly, we can show that a necessary and sufficient condition for w to commute with y is that $\epsilon a = \delta c$ whenever $\lambda(a, c) \neq 0$. Hence, since $\delta^2 = 1$, we can express the central element w in the form

$$w = \sum_{a \in \mathbb{Z}} \lambda(a) x^a z^{\epsilon \delta a} = \sum_{a \in \mathbb{Z}} \lambda(a) (w_{\delta, \epsilon})^{\delta a} \in \mathbb{K}[w_{\delta, \epsilon}^{\pm 1}].$$

Note that the integer powers of $w_{\delta, \epsilon}$ are linearly independent, and therefore the subalgebra of $\mathbb{K}_{\delta, \epsilon}[x^{\pm 1}, y^{\pm 1}, z^{\pm 1}]$ generated by $w_{\delta, \epsilon}$ and $w_{\delta, \epsilon}^{-1}$ is indeed a Laurent polynomial algebra. \square

It follows from the stratification theory of Goodearl and Letzter and the lemma above that the primitive ideals of $\mathbb{K}_{\delta,\epsilon}[x, y, z]$ with no nonzero homogeneous elements (relative to the grading induced by the isomorphisms of (27) and (28)) are those of the form

$$I_{\delta,\epsilon}^\alpha = \mathbb{K}_{\delta,\epsilon}[x, y, z] \cap \mathbb{K}_{\delta,\epsilon}[x^{\pm 1}, y^{\pm 1}, z^{\pm 1}](w_{\delta,\epsilon} - \alpha), \quad (30)$$

for $\alpha \in \mathbb{K}^*$.

Lemma 4.17. *Let $\alpha \in \mathbb{K}^*$. Then,*

- (a) $I_{1,1}^\alpha = \mathbb{K}_{1,1}[x, y, z](xz - \alpha)$;
- (b) $I_{-1,-1}^\alpha = \mathbb{K}_{-1,-1}[x, y, z](xz - \alpha^{-1})$;
- (c) $I_{1,-1}^\alpha = \mathbb{K}_{1,-1}[x, y, z](x - \alpha z)$;
- (d) $I_{-1,1}^\alpha = \mathbb{K}_{-1,1}[x, y, z](z - \alpha x)$.

Proof. We prove only (c), the other parts having similar proofs. Let us show the inclusion

$$I_{1,-1}^\alpha \subseteq \mathbb{K}_{1,-1}[x, y, z](x - \alpha z),$$

as the reverse one is clear. First notice that $x - \alpha z$ is a normal element of $\mathbb{K}_{1,-1}[x, y, z]$, and so $\mathbb{K}_{1,-1}[x, y, z](x - \alpha z)$ is indeed an ideal of $\mathbb{K}_{1,-1}[x, y, z]$. Assume

$$p = \left(\sum_{(a,b,c) \in \mathbb{Z}^3} \lambda(a, b, c) x^a y^b z^c \right) (xz^{-1} - \alpha) \in \mathbb{K}_{1,-1}[x, y, z], \quad (31)$$

where the sum in (31) is finite. Let $S = \{(a, b, c) \in \mathbb{Z}^3 \mid \lambda(a, b, c) \neq 0\}$ and suppose that there is $(a, b, c) \in S$ with $c \leq 0$. Define

$$c_0 = \min\{c \in \mathbb{Z} \mid \exists a, b \in \mathbb{Z} : (a, b, c) \in S\},$$

say $(a_0, b_0, c_0) \in S$. Then $c_0 - 1 < 0$, and since

$$p = \sum_{(a,b,c) \in S} \lambda(a, b, c) q^b x^{a+1} y^b z^{c-1} - \alpha \sum_{(a,b,c) \in S} \lambda(a, b, c) x^a y^b z^c, \quad (32)$$

the nonzero term $\lambda(a_0, b_0, c_0) q^{b_0} x^{a_0+1} y^{b_0} z^{c_0-1}$ which occurs in the sum (32), on the left, must be cancelled out by a term of the form $\alpha \lambda(a', b', c') x^{a'} y^{b'} z^{c'}$ occurring in the right-hand side sum of (32). This clearly contradicts the minimality of c_0 , as it would have to be $c' = c_0 - 1 < c_0$. Thus, $c \geq 1$ for all $(a, b, c) \in S$. One shows that $a, b \geq 0$ for all $(a, b, c) \in S$ in a similar fashion. Hence,

$$\sum_{(a,b,c) \in S} \lambda(a, b, c) x^a y^b z^{c-1} \in \mathbb{K}_{1,-1}[x, y, z] \quad (33)$$

and

$$p = \left(\sum_{(a,b,c) \in S} \lambda(a, b, c) x^a y^b z^{c-1} \right) (x - \alpha z), \quad (34)$$

as required. \square

Proposition 4.18. (a) $\text{Prim}_{J_{8,a}} U_q(\mathfrak{sl}_4^+) = \{(e_1e_2 - qe_2e_1, e_1e_3 - \alpha) \mid \alpha \in \mathbb{K}^*\};$

(b) $\text{Prim}_{J_{8,b}} U_q(\mathfrak{sl}_4^+) = \{(e_1e_2 - q^{-1}e_2e_1, e_1e_3 - \alpha) \mid \alpha \in \mathbb{K}^*\};$

(c) $\text{Prim}_{J_9} U_q(\mathfrak{sl}_4^+) = \{(e_1e_2 - q^{-1}e_2e_1, e_1 - \alpha e_3) \mid \alpha \in \mathbb{K}^*\};$

(d) $\text{Prim}_{J_{10}} U_q(\mathfrak{sl}_4^+) = \{(e_1e_2 - qe_2e_1, e_1 - \alpha e_3) \mid \alpha \in \mathbb{K}^*\}.$

The primitive ideals described above have height 4.

Proof. Parts (a)–(d) are a consequence of the isomorphisms of (27) and (28), Theorem 2.3, [10, Thm. 4.4] and Lemma 4.17, noting that for all $\alpha \in \mathbb{K}^*$

$$(e_1e_2 - qe_2e_1, e_2e_3 - qe_3e_2, e_1e_3 - \alpha) = (e_1e_2 - qe_2e_1, e_1e_3 - \alpha),$$

as two-sided ideals of $U_q(\mathfrak{sl}_4^+)$, and similarly for the other ideals involved in this proposition.

Let $P \in \text{Prim}_{J_{8,a}} U_q(\mathfrak{sl}_4^+)$, say $e_1e_3 - \alpha \in P$, with $\alpha \neq 0$. There is an algebra isomorphism $\mathbb{K}_q[\rho^{\pm 1}, \theta] \rightarrow U_q(\mathfrak{sl}_4^+)/P$ sending ρ, θ, ρ^{-1} to $e_1, e_2, \alpha^{-1}e_3$, respectively, where $\mathbb{K}_q[\rho^{\pm 1}, \theta]$ is the algebra generated by ρ, ρ^{-1} and θ , subject to the relations

$$\rho\rho^{-1} = 1 = \rho^{-1}\rho, \quad \rho\theta = q\theta\rho. \quad (35)$$

Since $\mathbb{K}_q[\rho^{\pm 1}, \theta]$ has GK dimension 2, it follows from Tauvel's height formula that P has height $6 - 2 = 4$. The other cases are analogous. \square

Consider the action of $U_q(\mathfrak{sl}_4^+)$ on the vector space $E = \mathbb{K}[t^{\pm 1}]$ of Laurent polynomials in the variable t , given by the formulas ($k \in \mathbb{Z}, \alpha \in \mathbb{K}^*$):

$$e_1.t^k = t^{k+1}, \quad e_2.t^k = q^k t^{k-1}, \quad e_3.t^k = \alpha t^{k-1}. \quad (36)$$

Then $E = E_\alpha$ becomes a simple $U_q(\mathfrak{sl}_4^+)$ -module and it is easy to see that

$$\text{ann } E_\alpha = (e_1e_2 - q^{-1}e_2e_1, e_1e_3 - \alpha).$$

To obtain simple $U_q(\mathfrak{sl}_4^+)$ -modules with annihilators in $\text{Prim}_{J_9} U_q(\mathfrak{sl}_4^+)$ we can consider the module E'_α , given by the following action of $U_q(\mathfrak{sl}_4^+)$ on $\mathbb{K}[t^{\pm 1}]$:

$$e_1.t^k = \alpha t^{k+1}, \quad e_2.t^k = q^k t^{k-1}, \quad e_3.t^k = t^{k+1}. \quad (37)$$

Finally, by twisting E_α by the automorphism η and changing q into q^{-1} in the formula for the action of e_2 on E'_α in (37), we get simple $U_q(\mathfrak{sl}_4^+)$ -modules with annihilators in $\text{Prim}_{J_{8,a}} U_q(\mathfrak{sl}_4^+)$ and $\text{Prim}_{J_{10}} U_q(\mathfrak{sl}_4^+)$, respectively.

4.3.9 The \mathcal{H} -strata of $(e_2e_3 - q^{\pm 1}e_3e_2, e_1)$ and $(e_1e_2 - q^{\pm 1}e_2e_1, e_3)$

Let $\mathbb{K}_q[x, y]$ be the *quantum plane*, given by generators x and y , satisfying only the relation $yx = qxy$. It is well-known that $\mathbb{K}_q[x, y]$ is primitive; in fact, let $\mathbb{K}_q[x, y]$ act on $\overline{F} = \mathbb{K}[t^{\pm 1}]$ as follows:

$$x.t^k = t^{k+1}, \quad y.t^k = q^k t^{k-1}, \quad k \in \mathbb{Z}.$$

Then, \overline{F} is a faithful representation of $\mathbb{K}_q[x, y]$ which is simple. Since there is an isomorphism

$$\mathbb{K}_q[x, y] \longrightarrow U_q(\mathfrak{sl}_4^+)/\langle e_2e_3 - qe_3e_2, e_1 \rangle, \quad (38)$$

mapping x to $e_3 + (e_2e_3 - qe_3e_2, e_1)$ and y to $e_2 + (e_2e_3 - qe_3e_2, e_1)$, we obtain the simple $U_q(\mathfrak{sl}_4^+)$ -module $F = \mathbb{K}[t^{\pm 1}]$ given by

$$e_1.t^k = 0, \quad e_2.t^k = q^k t^{k-1}, \quad e_3.t^k = t^{k+1}, \quad k \in \mathbb{Z}. \quad (39)$$

Note that $\text{ann } F = (e_2e_3 - qe_3e_2, e_1)$, by construction. Simple $U_q(\mathfrak{sl}_4^+)$ -modules with annihilators $(e_2e_3 - q^{-1}e_3e_2, e_1)$ and $(e_1e_2 - q^{\pm 1}e_2e_1, e_3)$ can be obtained, as before, by interchanging q and q^{-1} in (39) above and by using the automorphism η .

Proposition 4.19. (a) $\text{Prim}_{J_{11,a}} U_q(\mathfrak{sl}_4^+) = \{(e_2e_3 - qe_3e_2, e_1)\};$

(b) $\text{Prim}_{J_{11,b}} U_q(\mathfrak{sl}_4^+) = \{(e_1e_2 - q^{-1}e_2e_1, e_3)\};$

(c) $\text{Prim}_{J_{12,a}} U_q(\mathfrak{sl}_4^+) = \{(e_2e_3 - q^{-1}e_3e_2, e_1)\};$

(d) $\text{Prim}_{J_{12,b}} U_q(\mathfrak{sl}_4^+) = \{(e_1e_2 - qe_2e_1, e_3)\}.$

The primitive ideals described above have height 4.

Proof. For (a), we have just seen that $(e_2e_3 - qe_3e_2, e_1) \in \text{Prim}_{J_{11,a}} U_q(\mathfrak{sl}_4^+)$. Equality holds because $\text{Prim}_{J_{11,a}} U_q(\mathfrak{sl}_4^+)$ consists of a single \mathcal{H} -orbit. Since $\text{GKdim}(\mathbb{K}_q[x, y]) = 2$, the last statement of the proposition follows from the isomorphism of (38) and Tauvel's height formula. Parts (b)–(d) are analogous. \square

References

- [1] J. Alev and F. Dumas, *Sur le corps des fractions de certaines algèbres quantiques*, J. Algebra **170** (1994), no. 1, 229–265.
- [2] ———, *Rigidité des plongements des quotients primitifs minimaux de $U_q(\mathfrak{sl}(2))$ dans l'algèbre quantique de Weyl-Hayashi*, Nagoya Math. J. **143** (1996), 119–146.
- [3] N. Andruskiewitsch and F. Dumas, *On the automorphisms of $U_q^+(\mathfrak{g})$* , arXiv:math.QA/0301066.
- [4] J.L. Bueso, J. Gómez Torrecillas, F.J. Lobillo, and F.J. Castro, *An introduction to effective calculus in quantum groups*, Rings, Hopf Algebras, and Brauer Groups (Antwerp/Brussels, 1996), Lecture Notes in Pure and Appl. Math., vol. 197, Dekker, New York, 1998, pp. 55–83.
- [5] P. Caldero, *Générateurs du centre de $\check{U}_q(\mathfrak{sl}(N+1))$* , Bull. Sci. Math. **118** (1994), no. 2, 177–208.
- [6] ———, *Sur le centre de $U_q(\mathfrak{n}^+)$* , Beiträge Algebra Geom. **35** (1994), no. 1, 13–24, Festschrift on the occasion of the 65th birthday of Otto Krötenheerdt.
- [7] ———, *Étude des q -commutations dans l'algèbre $U_q(\mathfrak{n}^+)$* , J. Algebra **178** (1995), no. 2, 444–457.

- [8] J. Dixmier, *Enveloping Algebras*, Graduate Studies in Mathematics, vol. 11, American Mathematical Society, Providence, RI, 1996, Revised reprint of the 1977 translation.
- [9] K.R. Goodearl and T.H. Lenagan, *Catenarity in quantum algebras*, J. Pure Appl. Algebra **111** (1996), no. 1-3, 123–142.
- [10] K.R. Goodearl and E.S. Letzter, *The Dixmier-Moeglin equivalence in quantum coordinate rings and quantized Weyl algebras*, Trans. Amer. Math. Soc. **352** (2000), no. 3, 1381–1403.
- [11] K.R. Goodearl and R.B. Warfield, Jr., *An Introduction to Noncommutative Noetherian Rings*, London Mathematical Society Student Texts, vol. 16, Cambridge University Press, Cambridge, 1989.
- [12] M. Gorelik, *The prime and the primitive spectra of a quantum Bruhat cell translate*, J. Algebra **227** (2000), no. 1, 211–253.
- [13] T. Hayashi, *q -analogues of Clifford and Weyl algebras—spinor and oscillator representations of quantum enveloping algebras*, Comm. Math. Phys. **127** (1990), no. 1, 129–144.
- [14] E.E. Kirkman and L.W. Small, *q -analogs of harmonic oscillators and related rings*, Israel J. Math. **81** (1993), no. 1-2, 111–127. MR 94e:17019
- [15] S. Launois, *Primitive ideals and automorphism group of $U_q^+(\mathfrak{B}_2)$* , arXiv:math.RA/0412358.
- [16] S. Lopes, *Separation of variables for $U_q(\mathfrak{sl}_{n+1})^+$* , Canad. Math. Bull., to appear.
- [17] ———, *On the structure and representation theory of the quantized enveloping algebra $U_q(\mathfrak{g})^+$ for \mathfrak{g} semisimple*, Ph.D. thesis, University of Wisconsin-Madison, 2003.
- [18] M.P. Malliavin, *Algèbre d’Heisenberg quantique*, Bull. Sci. Math. **118** (1994), no. 6, 511–537.
- [19] J.C. McConnell and J.C. Robson, *Noncommutative Noetherian Rings*, revised ed., Graduate Studies in Mathematics, vol. 30, American Mathematical Society, Providence, RI, 2001, With the cooperation of L.W. Small.
- [20] C.M. Ringel, *PBW-bases of quantum groups*, J. Reine Angew. Math. **470** (1996), 51–88.