

Two issues in partially hyperbolic dynamics

Marcelo Viana
IMPA

based on joint work with A. Avila and A. Wilkinson

Partially hyperbolic dynamics

A diffeomorphism $f : M \rightarrow M$ on a compact Riemannian manifold is **partially hyperbolic** if there exists a continuous decomposition

$$T_x M = E_x^u \oplus E_x^c \oplus E_x^s$$

which is invariant under the dynamics:

$$Df_x(E_x^*) = E_{f(x)}^* \text{ for all } * \in \{u, c, s\}$$

and ...

- E^s is uniformly expanding:

$$\|Df_x|_{E_x^s}\| \leq \lambda < 1$$

- E^u is uniformly expanding:

$$\|(Df_x|_{E_x^u})^{-1}\| \leq \lambda < 1$$

- E^c is “in between”:

$$\frac{1}{\lambda} \frac{\|Df_x v^s\|}{\|v^s\|} \leq \frac{\|Df_x v^c\|}{\|v^c\|} \leq \lambda \frac{\|Df_x v^u\|}{\|v^u\|}$$

Uniformly hyperbolic dynamics

In the special case $E_x^c \equiv 0$, we say that f is **uniformly hyperbolic** (or **Anosov**), a notion that goes back to S. Smale, D. Anosov and Ya. Sinai in the 1960's.

Partial hyperbolicity is the most successful of the generalizations proposed in the 1970's, and has been a major topic in the dynamics over the last quarter of a century:

- they share many of the geometric properties of uniformly hyperbolic maps;
- they include many (new) interesting examples and phenomena;
- they are a suitable testing ground for outstanding issues in dynamics.

Many new examples

Basic fact: **partial hyperbolicity is an open property.**

- Take $A \in \mathrm{SL}(d, \mathbb{Z})$ whose spectrum intersects the interior, the boundary, and the exterior of the unit disk in \mathbb{C} . Then the **induced map** is partially hyperbolic:

$$f_A : \mathbb{T}^d \rightarrow \mathbb{T}^d, \quad \mathbb{T}^d = \mathbb{R}^d / \mathbb{Z}^d$$

Many new examples

Basic fact: **partial hyperbolicity is an open property.**

- Take $A \in \mathrm{SL}(d, \mathbb{Z})$ whose spectrum intersects the interior, the boundary, and the exterior of the unit disk in \mathbb{C} . Then the **induced map** is partially hyperbolic:

$$f_A : \mathbb{T}^d \rightarrow \mathbb{T}^d, \quad \mathbb{T}^d = \mathbb{R}^d / \mathbb{Z}^d$$

- Let $f^t : M \rightarrow M$, $t \in \mathbb{R}$ be an **Anosov flow**: there is an invariant decomposition

$$T_x M = E_x^u \oplus \mathbb{R}X(x) \oplus E_x^s, \quad X = \text{associated vector field.}$$

Then the **time-1 map** f^1 is partially hyperbolic.

Many new examples

Basic fact: **partial hyperbolicity is an open property.**

- Take $A \in \mathrm{SL}(d, \mathbb{Z})$ whose spectrum intersects the interior, the boundary, and the exterior of the unit disk in \mathbb{C} . Then the **induced map** is partially hyperbolic:

$$f_A : \mathbb{T}^d \rightarrow \mathbb{T}^d, \quad \mathbb{T}^d = \mathbb{R}^d / \mathbb{Z}^d$$

- Let $f^t : M \rightarrow M$, $t \in \mathbb{R}$ be an **Anosov flow**: there is an invariant decomposition

$$T_x M = E_x^u \oplus \mathbb{R}X(x) \oplus E_x^s, \quad X = \text{associated vector field.}$$

Then the **time-1 map** f^1 is partially hyperbolic.

- Let $g : N \rightarrow N$ be Anosov. Then any **isometry extension** is partially hyperbolic:

$$f : N \times \mathbb{T}^d \rightarrow N \times \mathbb{T}^d, \quad f(x, v) = (g(x), v + \omega(x)).$$

Invariant foliations: absolute continuity


Theorem (Anosov, Sinai, Brin, Pesin, Hirsch, Pugh, Shub)

Assume that $f : M \rightarrow M$ is partially hyperbolic. Then:

- The stable and unstable bundles are **uniquely integrable**: there exist unique foliations \mathcal{F}^s and \mathcal{F}^u such that

$$T_x \mathcal{F}_x^s = E_x^s \text{ and } T_x \mathcal{F}_x^u = E_x^u \text{ everywhere.}$$

- Those foliations \mathcal{F}^s and \mathcal{F}^u are **absolutely continuous**: projections along the leaves send zero measure sets to zero measure sets.



Absolute continuity of foliations (in the uniformly hyperbolic case) was the key ingredient in the proof of the famous result:

Theorem (Anosov)

The geodesic flow on a compact manifold with negative curvature is ergodic.

Absolute continuity of foliations (in the uniformly hyperbolic case) was the key ingredient in the proof of the famous result:

Theorem (Anosov)

The geodesic flow on a compact manifold with negative curvature is ergodic.

In contrast, a **center foliation** \mathcal{F}^c , tangent to E^c ,

- need not exist, nor be unique when it exists;
- need not be absolutely continuous.

A simple model

Consider the isometry extension $f_0 = g_A \times \text{id} : \mathbb{T}^2 \times \mathbb{T}^1 \rightarrow \mathbb{T}^2 \times \mathbb{T}^1$ of the map

$$g_A : \mathbb{T}^2 \rightarrow \mathbb{T}^2 \text{ is induced by } A = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}.$$

Known fact: Every diffeomorphism f of $\mathbb{T}^3 = \mathbb{T}^2 \times \mathbb{T}^1$ close to f_0 is partially hyperbolic with a unique center foliation, and the center leaves are smooth circles.

In what follows, always take $f : \mathbb{T}^3 \rightarrow \mathbb{T}^3$ to be volume preserving.

The (average) **center Lyapunov exponent** is the number

$$\lambda(f) = \int_{\mathbb{T}^3} \log |Df|_{E^c}.$$


Theorem (Shub, Wilkinson)

There are ergodic diffeomorphisms f close to f_0 such that $\lambda(f) \neq 0$.
Then the center foliation \mathcal{F}^c cannot be absolutely continuous.

Atomic disintegration

Theorem (Ruelle, Wilkinson)

If $\lambda(f) \neq 0$ then there exist $k \geq 1$ and a full volume subset of \mathbb{T}^3 that intersects every center leaf at exactly k points.



(there exists a full area subset of the square consisting of exactly 1 point on each of these curves)

Let $f : \mathbb{T}^3 \rightarrow \mathbb{T}^3$ be any C^k (volume preserving) diffeomorphism close to $f_0 : \mathbb{T}^3 \rightarrow \mathbb{T}^3$.

Theorem (Avila, Viana, Wilkinson)

If the center foliation \mathcal{F}^c is absolutely continuous, then f is C^k -conjugate to a rotation extension

$$\mathbb{T}^2 \times \mathbb{T}^1 \rightarrow \mathbb{T}^2 \times \mathbb{T}^1, \quad (x, v) \mapsto (g(x), v + \omega(x))$$

of some Anosov $g : \mathbb{T}^2 \rightarrow \mathbb{T}^2$, and the foliation \mathcal{F}^c is actually C^k .

Dichotomy theorem

Assume also that $f : \mathbb{T}^3 \rightarrow \mathbb{T}^3$ is **accessible**: any two points of \mathbb{T}^3 may be joined by a piecewise smooth path whose legs are contained in \mathcal{F}^s or \mathcal{F}^u leaves.

Theorem (Avila, Viana, Wilkinson)

Either the center foliation \mathcal{F}^c is absolutely continuous or there exists $k \geq 1$ and a full volume subset that intersects every center leaf at exactly k points. Generically, $k = 1$.

Accessibility is a mild assumption: it is known to be an open and dense in this setting (Nitičă, Török).

Let $f : T^1S \rightarrow T^1S$ be any diffeomorphism close to the time-1 map of the geodesic flow on a surface S with negative curvature.

Theorem (Avila, Viana, Wilkinson)

- 1 Either the foliation \mathcal{F}^c is C^∞ , or there exists some full volume subset that intersects every center leaf at exactly one point.
- 2 In the first case, f is the time-1 map of a C^∞ flow, whose trajectories coincide with the leaves of \mathcal{F}^c .

Let $f : T^1S \rightarrow T^1S$ be any diffeomorphism close to the time-1 map of the geodesic flow on a surface S with negative curvature.

Theorem (Avila, Viana, Wilkinson)

- 1 Either the foliation \mathcal{F}^c is C^∞ , or there exists some full volume subset that intersects every center leaf at exactly one point.
- 2 In the first case, f is the time-1 map of a C^∞ flow, whose trajectories coincide with the leaves of \mathcal{F}^c .

The proofs are based on a machinery (Invariance Principle etc) developed jointly with C. Bonatti, A. Avila and J. Santamaria, that is also used in the following result.

The **center Lyapunov exponents** of f are the numbers

$$\lambda(v) = \lim_n \frac{1}{n} \log \|Df_x^n(v^c)\| \text{ of vectors } v^c \in E_x^c$$

They are well defined almost everywhere (Oseledets theorem).

Question: **Can we always perturb f to make all the center Lyapunov exponents non-zero?**

Symplectic diffeomorphisms

Let $f_A : \mathbb{T}^4 \rightarrow \mathbb{T}^4$ be induced by some $A \in \mathrm{SL}(4, \mathbb{Z})$ with two eigenvalues in the unit circle.

Basic facts:

- f_A preserves some (constant) symplectic form ω .
- f_A preserves volume.
- Assuming that no eigenvalue is a root of unit, f_A is ergodic.

F. Rodriguez-Hertz: every volume preserving diffeomorphism f close to f_A is ergodic ([stable ergodicity](#)).

Symplectic diffeomorphisms

In fact, f is **stably Bernoulli** among symplectic diffeomorphisms:


Theorem (Avila, Viana)

Let $f : \mathbb{T}^4 \rightarrow \mathbb{T}^4$ be any ω -symplectic diffeomorphism close to f_A .

Then:

- either f has all center Lyapunov exponents non-zero,
- or f is conjugate to f_A by a volume preserving diffeomorphism.

In either case, f is ergodically equivalent to a Bernoulli shift.



4 issues/year

Electronic access

link.springer.com/journal/574/50/1

Subscription information

springer.com/librarians

Indexing

ZBMath, Science Citation Index
Scopus, MathSciNet, EBSCO,
OCLC WorldCat

Bulletin of the Brazilian Mathematical Society, New Series

Boletim da Sociedade Brasileira de Matemática

Editor-in-Chief: M. Viana

The Bulletin of the Brazilian Mathematical Society celebrates its 50th Volume in 2019. The Journal was relaunched in 1989 in a new edition with a new highly international editorial board. The Journal publishes novel research in mathematics of all fields, and forms a key part of the Brazilian Mathematical Society's mission.

Impact Factor: 0.410 (2017), Journal Citation Reports®

Throughout its history the Journal has published papers of importance from distinguished mathematicians working in all areas of mathematics. Some selected highlights from the Journal's history include:

Homomorphisms between Poisson JC-Algebras

Park*, CG. Bull Braz Math Soc, New Series (2005) 36: 79

A fixed point approach to stability of a quadratic equation

Mirzavaziri, M. & Moslehian, M.S. Bull Braz Math Soc, New Series (2006) 37: 361

The geometry of closed conformal vector fields on Riemannian spaces

Caminha, A. Bull Braz Math Soc, New Series (2011) 42: 277

On the homepage of **Bulletin of the Brazilian Mathematical Society, New Series** at **springer.com** you can

Sign up for our Table of Contents Alerts
Get to know the complete Editorial Board
Find submission information